

HOMEWORK 10, HONORS CALCULUS II
11/15/2018

Problem 1. Show that the generalized Newton binomial series (see previous HW sheet 9)

$$(1+x)^{1/2} = \sum_{k=0}^{\infty} \binom{\frac{1}{2}}{k} x^k$$

converges for all $x \in (-1, 1]$. Use this to calculate an approximate value for $\sqrt{2}$ by summing the first 5 terms of the series. How good is this approximation (compare to the value of $\sqrt{2}$ the calculator provides).

Problem 2. The curve $\gamma: [0, 2\pi] \rightarrow \mathbb{R}^2$ given by $\gamma(t) = (a \cos t, b \sin t)$ traces out an ellipse. We assume $0 < b < a$, so that a is the main axis.

- (i) Write down the integral which gives you a quarter of the length of the ellipse's circumference. This is called an *elliptic integral* and people have long tried to find an antiderivative without success. In fact, the antiderivative is a new type of function, called an *elliptic function*, which I believe was first properly understood by Riemann. So don't waste time trying to solve the integral by our methods.
- (ii) Putting the eccentricity $k = \frac{a^2 - b^2}{b^2}$ (so a circle has $k = 0$), you should have obtained $L = b \int_0^{\pi/2} \sqrt{1 + k(\sin t)^2} dt$ for the quarter length of the ellipse. Now use Newton's generalized binomial formula (see previous problem) to expand the integrand into a series and then integrate term by term. There will be a restriction on k for this series to converge (what is this restriction?).
- (iii) Choose $a = 2$ and $b = 3/2$ (then the series will converge, why?), and calculate an approximate quarter length L by summing the first 3 terms of your series expansion. Can you get a sense how the eccentricity influences the circumference of an ellipse?

Problem 3. Find a power series expansion for

$$\tan^{-1} x = \sum_{k=0}^{\infty} a_k x^k$$

Hint: $\int \frac{dx}{1+x^2} = \tan^{-1}(x) + c$ and expand the integrand into a series, integrate term by term, and determine what the constant c has to be. Find all x for which this series converges.

Problem 4. Determine *all* values for x such that the series

$$\sum_{k=1}^{\infty} \frac{x^k}{k}$$

converges.

Problem 5. Determine and provide proof of whether the following series of numbers converge or diverge:

- (i) $\sum_{n=1}^{\infty} \frac{n^{2n}}{(1+n)^{3n}}$
- (ii) $\sum_{k=1}^{\infty} (-1)^k \frac{k^4}{4^k}$
- (iii) $\sum_{k=1}^{\infty} \tan(1/k)$

- (iv) $\sum_{n=0}^{\infty} n^2 e^{-n^2}$
- (v) $\sum_{n=1}^{\infty} \frac{\sin n}{n\sqrt{n+1}}$
- (vi) $\sum_{n=1}^{\infty} \frac{n \ln n}{(n+1)^3}$
- (vii) $\sum_{n=1}^{\infty} \frac{(n!)^n}{n^{4n}}$
- (viii) $\sum_{n=1}^{\infty} n \sin(1/n)$
- (ix) $\sum_{n=1}^{\infty} (-1)^n \frac{\sqrt{n}}{n+3}$