

Math 461 Midterm review problems

Paul Hacking

October 21, 2018

- (1) A *rhombus* is a 4-sided polygon such that all the sides have the same length.
 - (a) Prove that the diagonals of a rhombus meet at right angles.
 - (b) Prove that the opposite sides of a rhombus are parallel.
- (2) Let $\triangle ABC$ be a triangle such that $\angle ACB = \pi/2$. Let L be the line through C perpendicular to AB . Let D be the intersection point of L and AB . Prove that $|AD| \cdot |DB| = |CD|^2$.

- (3) Let A, B, C, D be points on a circle and suppose that the chords AB and CD meet at a point P inside the circle. Prove that $|AP||BP| = |CP||DP|$.
- (4) Suppose given a circle \mathcal{C} . Give a ruler and compass construction of a rectangle $ABCD$ such that the vertices A, B, C, D lie on the circle and the angle between the diagonals is $\pi/6$.
[Note: For this problem and the other ruler and compass problems below, you may use the ruler and compass constructions described in class and in the text book (perpendicular bisector of a line segment, angle bisector, etc.) as components of your construction.]

- (5) Suppose given a circle \mathcal{C} . Give a ruler and compass construction of an isosceles right-angled triangle $\triangle ABC$ such that the sides are tangent to the circle.
- (6) Suppose given an angle θ and two lengths l and h . Give a ruler and compass construction of a triangle ABC with $\angle CAB = \theta$, $|BC| = l$, and perpendicular height h from A to BC . (Here we assume that $\tan(\theta/2) \leq l/2h$, otherwise such a triangle does not exist (why?).)

[Hint: First construct a triangle $A'BC$ with $\angle CA'B = \theta$ and $|BC| = l$. What is the locus of points A such that $\angle CAB = \theta$?

- (7) Find the intersection points of the circle with center the origin and radius 3 and the circle with center $(1, 2)$ and radius 2.
- (8) Find the center and radius of the circle passing through the points $A = (0, 0)$, $B = (1, 2)$, and $C = (-1, 3)$.
- (9) Show that the locus of points in the plane equidistant from a line L and a point P not on L is a parabola.
[Hint: Without loss of generality, we can choose coordinates such that the line L is the x -axis and the point P has coordinates $(0, b)$ for some $b \neq 0$.]
- (10) Suppose given a triangle $\triangle ABC$ such that $\angle ABC$ and $\angle BAC$ are acute

(less than $\pi/2$). What is the maximum area of a rectangle $DEFG$ with base DE on the side AB , vertex F on the side BC , and vertex G on the side CA ?

- (11) Let L be a line in \mathbb{R}^2 . Recall that Refl_L denotes reflection in the line L , an isometry of \mathbb{R}^2 .

Let P, Q be two points in \mathbb{R}^2 on the same side as L . Suppose we place a mirror on the line L (standing vertically, perpendicular to the plane), with the silvered side pointing towards P and Q . Show that the perceived position of P for an observer at Q looking in the mirror is $\text{Refl}_L(P)$. (In other words, light from the object at P reflected in the mirror travels the

same distance to Q and arrives in the same direction as light from an object at $\text{Refl}_L(P)$ when the mirror is removed.)

[Hint: For reflection of light rays in a mirror, the incoming ray makes the same angle with the mirror as the outgoing ray (“angle of incidence equals angle of reflection”).]

(12) Give a precise geometric description for each of the following isometries $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ as a translation, rotation, reflection, or glide reflection.

(a) $T(x, y) = (x + 5, y + 7)$.

(b) $T(x, y) = (1 - y, 3 - x)$.

(c) $T(x, y) = (y + 2, 8 - x)$.

(d) $T(x, y) = \frac{1}{13}(5x + 12y + 4, 12x -$

$5y - 6$).

- (13) Give an algebraic formula for the rotation about the point $(4, 2)$ through angle $\pi/4$ radians counterclockwise.
- (14)(a) Let L be a line through the origin in \mathbb{R}^2 . Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the reflection in L . Let \mathbf{n} be a vector perpendicular to L . Show that T is given by the formula

$$T(\mathbf{x}) = \mathbf{x} - 2 \left(\frac{\mathbf{x} \cdot \mathbf{n}}{\mathbf{n} \cdot \mathbf{n}} \right) \mathbf{n}.$$

[Hint: Here $\mathbf{u} \cdot \mathbf{v}$ is the dot product, defined by $(u_1, u_2) \cdot (v_1, v_2) = u_1v_1 + u_2v_2$. In particular $\mathbf{u} \cdot \mathbf{u} = \|\mathbf{u}\|^2$, where $\|\mathbf{u}\|$ is the length of the vector \mathbf{u} . Recall that we have the formula

$$\mathbf{u} \cdot \mathbf{v} = \|\mathbf{u}\| \|\mathbf{v}\| \cos \theta$$

where θ is the angle between the vectors \mathbf{u} and \mathbf{v} .]

- (b) Find an algebraic formula for the reflection in the line $y = 3x$.
 - (c) Find an algebraic formula for the reflection in the line $y = 3x + 1$.
- (15) Give a precise geometric description for each of the following isometries as a translation, rotation, reflection, or glide reflection.
- (a) The composition of reflection in the line $x = 3$ followed by reflection in the line $y = \sqrt{3}x$.
 - (b) The composition of reflection in the line $y = 2x + 1$ followed by reflection in the line $y = 2x + 5$.
 - (c) The composition of rotation about

the point $(1, 1)$ through angle $\pi/2$ counterclockwise followed by rotation about the point $(2, 2)$ through angle $\pi/2$ counterclockwise.

(d) The composition of rotation about the point $(3, 2)$ through angle $\pi/3$ counterclockwise followed by translation by $(4, 0)$

(16) Let L and M be two distinct lines in the plane. Show that $\text{Refl}_M \circ \text{Refl}_L = \text{Refl}_L \circ \text{Refl}_M$ iff L and M are perpendicular.

(17) Recall that $\text{Rot}(P, \theta)$ denotes rotation about a point P through angle θ counterclockwise. Let $ABCD$ be a square such that the vertices A, B, C, D are in counterclockwise order. Determine the composition

$\text{Rot}(A, \pi/2) \circ \text{Rot}(B, \pi/2) \circ \text{Rot}(C, \pi/2) \circ \text{Rot}(D, \pi/2)$.

Justify your answer carefully.

- (18) Express the isometry $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(x, y) = (x+6, 4-y)$ as a composite of at most 3 reflections.
- (19) Let $A = (0, 0)$, $B = (1, 0)$, $C = (0, 2)$, and $A' = (3, 5)$, $B' = (3, 4)$, $C' = (1, 5)$. Find an isometry T sending $\triangle ABC$ to $\triangle A'B'C'$. (Give a geometric description and an explicit algebraic formula.)