

The GHS Inequality for a Large External Field

Richard S. Ellis,^{1,3} Charles M. Newman,^{2,4} and Michael R. O'Connell¹

Received November 5, 1980

We consider general even ferromagnetic systems with pair interactions in a nonnegative external magnetic field h . Classes of single-site measures ρ are found such that the GHS inequality is valid for all $h \geq \tilde{h}$, where $\tilde{h} \geq 0$ is a number depending on ρ but independent of the size of the system. These measures include both absolutely continuous and discrete measures. For $\rho \doteq a\delta_0 + \{(1-a)/2\} \cdot (\delta_1 + \delta_{-1})$, some $a \in [0, 1)$, \tilde{h} is determined exactly.

KEY WORDS: GHS inequality; general even ferromagnetic systems; correlation inequalities.

1. INTRODUCTION

The Griffiths–Hurst–Sherman (GHS) inequality is a useful tool in the study of lattice spin systems with ferromagnetic pair interactions. For example, when valid, it implies that the average magnetization per site is a concave function of $h \geq 0$, where h denotes the external magnetic field. It also implies the absence of spontaneous magnetization except possibly at $h = 0$.⁽⁹⁾ However, the validity of the GHS inequality for a particular system depends upon that system's single spin measures. For example, it holds for spin-1/2 systems⁽⁵⁾—i.e., for systems with single spin measures the Bernoulli measure $\frac{1}{2}(\delta_1 + \delta_{-1})$ —but not for systems with single spin

¹ Department of Mathematics and Statistics, University of Massachusetts, Amherst, Massachusetts 01003.

² Department of Mathematics, University of Arizona, Tucson, Arizona 85721.

³ Alfred P. Sloan Research Fellow. Research supported in part by National Science Foundation grant No. MCS 80-02149.

⁴ Alfred P. Sloan Research Fellow. Research supported in part by National Science Foundation grant No. MCS 77-20683 and by the U.S.–Israel Binational Science Foundation.

