

Lecture Notes in Mathematics

Edited by A. Dold and B. Eckmann

451

Probabilistic Methods in Differential Equations

Proceedings of the Conference Held at the
University of Victoria, August 19–20, 1974

Edited by M. A. Pinsky



Springer-Verlag
Berlin · Heidelberg · New York 1975

Asymptotics and Limit Theorems
for the Linearized Boltzmann Equation¹

Richard S. Ellis²

We consider the initial value problem for the linearized Boltzmann equation

$$(1) \quad \frac{\partial p}{\partial t} + \xi \cdot \text{grad } p = \frac{1}{\epsilon} Qp, \quad \lim_{t \downarrow 0} p = f,$$

where the initial data $f = f(x, \xi)$ and the solution $p = p_\epsilon(t, x, \xi)$, $t > 0$, $x \in \mathbb{R}^3$, $\xi = (\xi_1, \xi_2, \xi_3) \in \mathbb{R}^3$. Q is the linearized collision operator corresponding to a spherically symmetric intermolecular potential, and $\epsilon > 0$ is a parameter which represents the mean free path. Corresponding to the conservation of number, momentum, and energy in an individual collision, Q has a five-dimensional nullspace spanned by 1, ξ_i ($i = 1, 2, 3$), and $|\xi|^2$.

We are interested in the asymptotic behavior of the solution of (1) as $\epsilon \downarrow 0$. This is formally treated at the physical level of rigor by the Chapman-Enskog-Hilbert procedure [9; pp. 254-262]. Given a nice f , we define

1. This is a report on joint work with Mark A. Pinsky.
2. Supported in part by National Science Foundation Grant GP-28576

$$f_0(x) \equiv \langle f(x, \cdot), 1 \rangle; f_i(x) \equiv \langle f(x, \cdot), \xi_i \rangle, i = 1, 2, 3;$$

$$(2) \quad f_4(x) \equiv \langle f(x, \cdot), \frac{|\xi|^2 - 3}{\sqrt{6}} \rangle;$$

$$n_0(t, x) \equiv \langle p_\epsilon(t, x, \cdot), 1 \rangle; n_i(t, x) \equiv \langle p_\epsilon(t, x, \cdot), \xi_i \rangle, i = 1, 2, 3;$$

$$n_4(t, x) \equiv \langle p_\epsilon(t, x, \cdot), \frac{|\xi|^2 - 3}{\sqrt{6}} \rangle,$$

where p_ϵ is the solution of (1) with initial data f and where $\langle \cdot, \cdot \rangle$ denotes the inner product in the Hilbert space

$\mathcal{H}_0 = L^2(\mathbb{R}^3; (2\pi)^{-3/2} \exp(-\frac{1}{2}|\xi|^2) d\xi)$. Provided f satisfies certain algebraic compatibility conditions (the Hilbert relations), one can show that up to an error $O(\epsilon)$ the n_i satisfy the following system of partial differential equations (linear Euler equations):

$$\frac{\partial n_0}{\partial t} = - \operatorname{div} \vec{n},$$

$$(4) \quad \frac{\partial \vec{n}}{\partial t} = - \operatorname{grad} n_0 - \sqrt{\frac{2}{3}} \operatorname{grad} n_4,$$

$$\frac{\partial n_4}{\partial t} = - \sqrt{\frac{2}{3}} \operatorname{div} \vec{n},$$

$$n_i(0^+, x) = f_i(x),$$

where $\vec{n} = (n_1, n_2, n_3)$. Further, if one writes $p_\epsilon(t/\epsilon, x, \cdot)$ for $p_\epsilon(t, x, \cdot)$ in (2), then up to an error $O(\epsilon)$ the resulting $n_i(\frac{t}{\epsilon}, x)$ agree with the solutions (evaluated at time t/ϵ) of the following system of partial differential equations (linear Navier-Stokes

