

SYMPLECTOMORPHISMS OF MIRRORS TO LOG CALABI-YAU SURFACES

PAUL HACKING AND AILSA KEATING

ABSTRACT. Let M be a Weinstein four-manifold mirror to $Y \setminus D$ for (Y, D) a log Calabi–Yau surface; intuitively, this is typically the Milnor fibre of a smoothing of a cusp singularity. We introduce two families of symplectomorphisms of M : Lagrangian translations, which we prove are mirror to tensors with line bundles; and nodal slide recombinations, which we prove are mirror to automorphisms of (Y, D) . The proof uses a detailed compatibility between the homological and SYZ view-points on mirror symmetry. Together with spherical twists, these symplectomorphisms are expected to generate all autoequivalences of the wrapped Fukaya category of M which are compactly supported in a categorical sense. A range of applications is given.

CONTENTS

1. Introduction	1
2. Autoequivalences of $D^b \text{Coh}(Y)$ and $D^b \text{Coh}(U)$	11
3. Mirror symmetry for log Calabi–Yau surfaces: background	19
4. Lagrangian translations and tensors by line bundles	25
5. Lagrangian spheres mirror to (-2) curves	34
6. Nodal slide recombinations	37
7. Applications	47
References	54

1. INTRODUCTION

1.1. **Context and main theorem.** Consider a symplectic manifold M , possibly with boundary. Let $\text{Symp}_c(M)$ be its group of compactly supported symplectomorphisms; our convention is that if $\partial M \neq \emptyset$, an element of $\text{Symp}_c(M)$ must be the identity on a collar neighbourhood of ∂M . We are interested in the symplectic mapping class group $\pi_0 \text{Symp}_c(M)$.

A rich source of mapping class elements are Dehn twists in Lagrangian spheres. First introduced by Arnol'd [Arn95] as the monodromy of complex Morse singularities, they have since been

extensively studied. Key to this was understanding their action on Fukaya categories: spherical twists autoequivalences [Sei03, Sei08]. On the B side of mirror symmetry, spherical twists acts on suitable derived categories of coherent sheaves [ST01]. Dehn twists are currently our best source of symplectic mapping class group elements, with generalisations such as fibred twists and twists in other Lagrangians with periodic geodesic flow [Sei00, Per, WW16]. In the small number of cases for which we have full descriptions of symplectic mapping class groups, they are generated by global symmetries (typically for closed manifolds) and Dehn twists. In particular, restricting to the case where M is a Weinstein domain, symplectic mapping class groups are known for (real) four-dimensional Milnor fibres of A_n singularities [KS02, Eva11, LW12]: we get the braid group on n strands.

In this paper, we introduce some new families of symplectic mapping classes of four-dimensional Weinstein domains.

Heuristically, the Weinstein domains M we work with are Milnor fibres of smoothings of simple elliptic or cusp singularities. They are mirror to maximally degenerate log Calabi–Yau surfaces (Y, D) , with the distinguished complex structure within their deformation class (following the notation in [GHK15b], $(Y, D) = (Y_e, D)$; this corresponds to the trivial period point). These singularities are typically of neither hypersurface nor complete intersection type. There is a conjectural description for the smoothing components of these singularities, see Section 1.5. From this description, one expects monodromy to give symplectomorphisms of M beyond Dehn twists.

We construct these symplectomorphisms directly, building on [GHK15b, HK]. There are two families: compactly supported symplectomorphisms which are mirror (in a sense which we will make precise) to tensors with line bundles; and to automorphisms of (Y, D) . Moreover, we will see that for such Weinstein domains M , we expect that together with Dehn twists, our symplectomorphisms generate the symplectic mapping class group of M , or rather its categorical representation as autoequivalences of the wrapped Fukaya category $\mathcal{W}(M)$. (The ambiguity can be pinned down: loosely, one needs to know that there cannot be exotic autoequivalences of the mirror with support on finite unions of (-2) curves, see Section 1.4.)

Remark 1.1. Several works already hinted at a rich story for symplectomorphisms beyond Dehn twists. In particular, Sheridan–Smith proved that Dehn twists can’t generate $\pi_0 \text{Symp}_c(M)$ for some K3 surfaces [SS20a], and Shevchishin–Smirnov constructed a four-dimensional ‘elliptic twist’ in the presence of a self-intersection (-1) symplectic two-torus [SS20b].

Mirror symmetry setting. We focus on the (complex) two-dimensional, non-compact Calabi–Yau setting. On the B side, (Y, D) is a log Calabi–Yau surface with maximal boundary and distinguished complex structure. Its mirror is a four-dimensional Weinstein domain M together with a superpotential $w : M \rightarrow \mathbb{C}$ [Kea18, HK]. Let U denote $Y \setminus D$. Homological mirror symmetry consists of a triple of compatible explicit equivalences, which we take to be fixed for

the rest of this section:

$$\mathrm{Perf}(D) \simeq D^\pi \mathcal{F}(\Sigma) \quad D(Y) \simeq D^b \mathcal{F}(w) \quad D(U) \simeq D^b \mathcal{W}(M)$$

where $D^b \mathcal{F}(w)$ is the derived directed Fukaya category of w , $D^b \mathcal{W}(M)$ the derived wrapped Fukaya category of M , and $D^\pi \mathcal{F}(\Sigma)$ the split-closed derived Fukaya category of a smooth fibre of w near infinity, see [HK, Theorem 1.1]. Typically, D is the cycle of a cusp singularity, and M should be thought of as the Milnor fibre of a smoothing of the dual cusp [Loo81, GHK15a, Eng18]. M can be described explicitly as a Weinstein handlebody, by starting with D^*T^2 and gluing on Weinstein two handles along the conormal Legendrian lifts of rational slope circles in T^2 , see [HK]. This is a rich family of examples, with connections to cluster varieties cf. [GHKK18, STW].

Categorical compact support. Suppose $f \in \pi_0 \mathrm{Samp}_c M$ is a mapping class element mirror to an autoequivalence $\check{f} \in \mathrm{Auteq} D(Y)$. (We will see F can be given a preferred grading.) As f fixes the fibres of w outside a compact set, homological mirror symmetry implies that \check{f} respects restriction to $\mathrm{Perf} D$, i.e. $\iota_D^* \circ \check{f} = \iota_D^*$. By abuse of terminology, we say that an autoequivalence of $D(Y)$ satisfying this condition has ‘compact support on $Y \setminus D$ ’.¹

Main theorem. Our main theorem is as follows.

Theorem 1.2. *Let (Y, D) be a log Calabi-Yau surface with distinguished complex structure, and $w : M \rightarrow \mathbb{C}$ its mirror.*

- (1) (Theorem 5.5.) *Suppose $\mathcal{L} \in \mathrm{Pic}(Y)$ is a line bundle such that $\mathcal{L}|_D = \mathcal{O}_D$. Then there exists a compactly supported symplectomorphism $\sigma_{\mathcal{L}}$ of M such that under homological mirror symmetry, $\otimes \mathcal{L} \in \mathrm{Auteq} D(U)$ is taken to $\sigma_{\mathcal{L}} \in \mathrm{Auteq} D^b \mathcal{W}(M)$. We call $\sigma_{\mathcal{L}}$ a Lagrangian translation.*
- (2) (Theorem 6.1.) *Suppose $\varphi \in \mathrm{Aut}(Y)$ is a biholomorphic automorphism fixing D pointwise. Then there exists a compactly supported symplectomorphism $\check{\varphi}$ of M such that under homological mirror symmetry, $\varphi_* \in \mathrm{Auteq} D(Y)$, respectively $\mathrm{Auteq} D(U)$, is taken to $\check{\varphi} \in \mathrm{Auteq} D^b \mathcal{F}(w)$, respectively $\mathrm{Auteq} D^b \mathcal{W}(M)$. We call $\check{\varphi}$ a nodal slide recombination.*

The names will be self-explanatory from the constructions, described below. Both families on the B side are understood: the line bundles in (1) lie in $Q = \langle D_1 \dots, D_k \rangle^\perp \subset \mathrm{Pic}(Y)$, where the D_i are the irreducible components of D . (This uses $(Y, D) = (Y_e, D)$.) We construct a symplectomorphism for each element of \bar{Q} , the image of Q in $\mathrm{Pic}(U) = \mathrm{Pic}(Y)/\langle D \rangle$, which satisfies $\bar{Q} \simeq H_2(U)/H_2(\partial U)$. For (2), the Torelli theorem in [GHK15b] implies that the group of automorphisms fixing D pointwise is identified with Adm/W in Gross–Hacking–Keel’s notation, where Adm is the group so-called admissible lattice automorphisms of $\mathrm{Pic}(Y)$, and

¹From the perspective of the geometry of Y , ‘asymptotically compactly supported’ would be more accurate.

W is the Weyl group. (See [GHK15b, Definition 4.2].) Geometrically, our constructions are well-behaved with respect to composition: we get a faithful group homomorphism (Theorem 7.1 part (1)):

$$\bar{Q} \rtimes \text{Adm}/W \rightarrow \pi_0 \text{Symp}_c(M).$$

Let $\langle \Phi \rangle \subset \bar{Q}$ be the sublattice generated by classes of (-2) curves in $Y \setminus D$. For any $\mathcal{L} \in \Phi$, we'll see that the symplectomorphism $\sigma_{\mathcal{L}}$ can be factorised as the product of two Dehn twists in Lagrangian spheres lifting these roots (Lemma 6.13). All other symplectomorphisms we construct are new, in the following sense. Denote by $\text{Br} \subset \pi_0 \text{Symp}_c M$ the subgroup generated by all Dehn twists, and by \bar{Q}_{symp} , etc, the images of our groups in $\pi_0 \text{Symp}_c(M)$. Theorem 7.1 part (2) tells us that:

$$(\bar{Q} \rtimes \text{Adm}/W)_{\text{symp}} \cap \text{Br} = \langle \Phi \rangle_{\text{symp}} \rtimes \{1\} \subset \pi_0 \text{Symp}_c M.$$

One of the first examples is the following.

Example 1.3. (Examples 6.11 and 7.4.) Let $X = \mathbb{F}_1$ be the blow-up of \mathbb{P}^2 at a point, and $M = X \setminus E$ the complement of a smooth anticanonical divisor. This is mirror to a log Calabi-Yau (Y, D) such that D is a cycle of eight (-2) curves. Then $\bar{Q} \simeq \mathbb{Z}$, $\Phi = \emptyset$, and $\text{Adm}/W \simeq \mathbb{Z}$. (The latter consists of eighth powers of elements of the Mordell-Weil group of Y , see Section 2.2.2.) This means we have a family of symplectomorphisms

$$\mathbb{Z}^2 \leq \pi_0 \text{Symp}_c(M).$$

Following [Gro85], $\text{Symp}(X, \omega_{FS})$ is homotopy equivalent to $U(2)$, where ω_{FS} is the Fubini–Study form. In particular, its symplectic mapping class group is trivial. On the other hand, we'll see that at the categorical level, \mathbb{Z}^2 gives all compactly supported symplectomorphisms of M . (We'll see that the boundary twist acts trivially on the wrapped Fukaya category $\mathcal{W}(M)$.)

A further range of examples is given in Section 7.1. Note that elements of Adm/W persist under blow-ups at points of D , which are precisely the ones which respect the log CY property; from that perspective, non-triviality of Adm/W should be thought of as generic.

1.2. Constructions. Both constructions are most natural from the perspective of an almost-toric fibration $\pi : M \rightarrow \mathbb{R}^2$ coming from the Gross–Siebert programme, which should be thought of as the SYZ fibration on M . Formally, to describe this, pick a toric model for (Y, D) : possibly after blowing up corners of D , we can find a toric pair (\bar{Y}, \bar{D}) and a map $(Y, D) \rightarrow (\bar{Y}, \bar{D})$ given by blowing up interior points of the \bar{D}_i . As Y has the distinguished complex structure, we blow up a single favourite point on \bar{D}_i , say m_i times. Let v_i be the corresponding ray in the fan for \bar{Y} . The associated almost-toric fibration π has m_i nodal fibres with invariant direction v_i , with all invariant lines concurrent. (See [HK, Theorem 1.2].) Changes of toric models give equivalent fibrations, which will be further exploited below.

Lagrangian translations. Fix a reference Lagrangian section L_0 for the almost-toric fibration $\pi : M \rightarrow \mathbb{R}^2$. Heuristically, this should be thought of as mirror to \mathcal{O}_Y . More generally, Lagrangian sections of π , up to suitable equivalence, are expected to correspond to line bundles on Y . Restricting to ‘compactly supported’ subsets on both sides, we will see in Corollary 4.20 that there are one-to-one correspondences

$$\left\{ \begin{array}{l} \text{Lagrangian sections } L \text{ of } \pi \text{ equal to } L_0 \text{ near } \partial M \\ \text{up to fibre-preserving Hamiltonian isotopy} \end{array} \right\} \longleftrightarrow H_2(M)/H_2(\partial M)$$

$$\begin{array}{ccc} & & \updownarrow \\ \text{Line bundles in } \bar{Q} & \longleftrightarrow & H_2(U)/H_2(\partial U) \end{array}$$

where the vertical equivalence is induced by an explicit diffeomorphism between M and U , the top one by taking L to $[L] - [L_0]$, and the bottom one by taking the first Chern class and Poincaré duality. To define a Lagrangian translation, fix a Lagrangian section L . Arnol’d–Liouville coordinates give a linear structure on each smooth fibre, and adding $L - L_0$ fibrewise is locally a symplectomorphism. We show that this can be extended over the singular fibres to get an element of $\pi_0 \text{Symp}_c(M)$ (Proposition 4.9).

This construction will seem natural to experts. In particular, it closely ties back to ideas of Gross for compact Calabi–Yau manifolds [Gro98]; and related constructions can be found, inter alia, in Subotic’s thesis [Sub10, Chap. 5], Hanlon–Hicks’ definition of the fibrewise sum of two Lagrangian sections [HH, Section 5.2], and work of Engel–Friedman [EF21] (see also Remark 1.11).

Nodal slide recombinations. Suppose φ is an automorphism of Y preserving D pointwise. To construct the mirror $\tilde{\varphi}$, we use input from birational geometry. Fix any toric model for (Y, D) , say $f : (Y, D) \rightarrow (\bar{Y}, \bar{D})$. (We can without loss of generality ignore toric blow-ups.) It follows from the two-dimensional Sarkisov programme that the toric model $f \circ \varphi : (Y, D) \rightarrow (\bar{Y}, \bar{D})$ can be obtained by starting from $f : (Y, D) \rightarrow (\bar{Y}, \bar{D})$ and performing a sequence of elementary transformations (see Proposition 3.5 and [CK16, KSC04]). Equivalently, φ restricts to a birational transformation of $(\mathbb{C}^*)^2$, say $\tilde{\varphi}$, and the sequence of elementary transformations corresponds to a factorisation of $\tilde{\varphi}$ into a composition of maps of the form $\psi \circ E \circ \psi^{-1}$, where $E : (x, y) \mapsto (x, y/(x + 1))$ and $\psi \in SL_2(\mathbb{Z})$ (cf. [Bla13]). Readers will recognise E as the standard two-variable cluster transformation.

On the mirror side, each elementary transformation determines an operation of M . In terms of the almost-toric fibration, this is a nodal slide followed by a cut transfer ([Sym03], see Section 3.2.2); in terms of the Weinstein skeleton of M , a Lagrangian mutation (for which we will follow [PT20]). The full sequence of transformations returns the same almost-toric fibration (or presentation of the Weinstein skeleton) as we started with, and determines a symplectomorphism of M . See Theorem 6.1. Independence of the choice of factorisation reduces to checking a symplectic incarnation of the ‘ A_2 cluster relation’, see Proposition 6.6.

This uses Blanc’s description of the relations in the group of volume-preserving birational transformations of \mathbb{P}^2 [Bla13].

Some explicit examples are calculated in Section 6.3.

The idea that elementary transformations are mirror to Lagrangian mutations has been widely explored in the wake of [Aur07] – see for instance [Via14, Via17, STW, PT20]. From the cluster perspective, the symplectomorphisms are interesting in of their own right: they are associated to loops in the cluster graph for $Y \setminus D$. For K3 surfaces, similar monodromy transformations were studied by Kontsevich–Soibelman in terms of the bases of the SYZ fibrations. See [KS06, Conjecture 7].

Heuristically, we expect a symplectomorphism to respect the SYZ fibration if and only if the mirror autoequivalence preserves the collection of structure sheaves of points. On the other hand, by [Huy06, Corollary 5.23], such autoequivalences are precisely given by $\text{Pic}(Y) \rtimes \text{Aut}(Y)$: this would mean that we have found all compactly supported symplectomorphisms respecting the SYZ fibration, up to equivalence in $\text{Auteq } \mathcal{W}(M)$.

Remark 1.4. The reader may be interested to note that from the perspective of mirror symmetry, the factorisation of $\bar{\varphi} \in \text{Bir}((\mathbb{C}^*)^2, \Omega)$ into elementary transformations seems more natural than the one into Cremona transformations, more commonly considered.

1.3. Homological mirror symmetry. In order to prove that the symplectomorphisms we construct have the expected mirrors, the main ingredients are as follows:

- (1) Precise geometric translations between the Lefschetz and SYZ fibrations on M , and operations on them, building on [HK]. This should be thought of as a fine-grain compatibility between the homological and SYZ perspectives for mirror symmetry.

This suffices for nodal slide recombinations. For Lagrangian translations, we also use the following:

- (2) Explicit constructions of Lagrangian spheres mirror to $i_*\mathcal{O}_C(a)$, for any $C \subset U$ a (-2) curve, and $a \in \mathbb{Z}$. These and their twists in themselves are all the known sphericals in $D(U)$.
- (3) Work of Uehara on autoequivalences of $D(Y)$ [Ueh16, Ueh19]. The key technical point is that whenever D is negative definite or indefinite, the Fourier–Mukai kernel of any autoequivalence of $D(Y)$ has two-dimensional support. (In the semi-definite case, we will instead exploit the fact that Y is rational elliptic.)

Combining (2) and (3), we can prove tractable criteria for an autoequivalence to be the identity: see for instance Proposition 2.14 or the proof of Theorem 5.5.

Non-compactly supported maps. Our constructions naturally extend to give examples of non-compactly supported symplectomorphisms:

- for Lagrangian translations, we can use a Lagrangian section L which does not agree with L_0 near infinity; see Lemma 4.3 and Proposition 4.12. We expect this to be mirror to $\otimes \mathcal{L}$ for a line bundle \mathcal{L} such that $\mathcal{L}|_D \not\cong \mathcal{O}_D$.
- for nodal slide recombinations, we can add extra automorphisms of Y , which fix D setwise but not pointwise, see Section 6.2.1. This corresponds to adding actions of elements of $SL_2(\mathbb{Z})$ on the base of the almost-toric fibration. We expect to get an action by symplectomorphisms of the cluster modular group introduced in [FG09].

For the latter, we can also describe the symplectomorphisms in terms of the Lefschetz fibration $w : M \rightarrow \mathbb{C}$, and use this to show that they act as expected on Fukaya categories, see Remark 6.5. For the former, it is natural to ask for mirrors to arbitrary elements of $\text{Pic}(Y)$; these should correspond to Lagrangian translations in sections with prescribed asymptotics near infinity. As such sections are studied in detail in work in preparation by Hanlon–Ward [HW] on HMS for log CY pairs via monomially admissible sections (building on [Han19, HH]), we do not pursue this here.

More generally, we expect that the monomially admissible framework, once established, will give us stronger technical tools for studying properties of our symplectomorphisms, including non-compactly supported ones: these maps are described most naturally in terms of the SYZ fibrations, and working with versions of the Fukaya–Seidel category defined directly in terms of these fibrations should allow us, for instance, to by-pass descriptions in terms of Lefschetz fibrations. Some properties of non-compactly supported maps will be studied in [HKW].

1.4. Further applications.

Autoequivalence groups. A full description of $\text{Auteq } D(Y)$ would require classifying all spherical objects in $D(Y)$.

Conjecture 1.5. *Let \mathcal{A} be the collection of spherical objects*

$$\mathcal{A} := \{i_* \mathcal{O}_C(a) \mid C \text{ a } (-2) \text{ curve, } a \in \mathbb{Z}\}.$$

Let $T_{\mathcal{A}}$ be the group generated by spherical twists in objects of \mathcal{A} . Then any spherical object in $D(Y)$ is quasi-isomorphic to an element of $T_{\mathcal{A}} \cdot \mathcal{A}$.

Any spherical object in $D(Y)$ has support in Z , the union of all (-2) curves in Y , and the conjecture is known for all connected components of Z which are either chains or cycles of (-2) curves ([IU05, Ueh16], see Lemma 2.1; note the extra technical assumption in the case where the cycle of (-2) curves is D itself). A companion question is to understand the full group of autoequivalences of $D(Y)$. For compactly supported ones, the general expectation translates as follows.

Conjecture 1.6. *Cf. [Ueh19, Conjecture 1.2]. Let $\text{Auteq}_c D(U)$ denote compactly supported autoequivalences of $D(U)$: restrictions of autoequivalences of $D(Y)$ which commute with the restriction to $\text{Perf}(D)$, see Section 2.4. Then*

$$\text{Auteq}_c D(U) = \langle T_{\mathcal{A}'}, \bar{Q} \rangle \rtimes \text{Aut}(Y, D; pt)$$

where $\mathcal{A}' \subset \mathcal{A}$ is given by restricting to curves in $Y \setminus D$, \bar{Q} is as before, and $\text{Aut}(Y, D; pt)$ is the group of automorphisms of Y fixing D pointwise.

This too is known to hold whenever all connected components of $Z \cap Y \setminus D$ are chains or cycles of (-2) curves [IU05, Ueh16, Ueh19]. Examples include the cases where D is semi-definite case and $k \geq 5$ (assuming D contains no (-1) curves). The difficulty is classifying autoequivalences of $D_Z(Y)$, the derived category of coherent sheaves with support on Z . While this problem is largely orthogonal to the focus of our article, note that our results can be packaged as follows:

Theorem 1.7. *(Proposition 5.2 and Theorem 7.1) Assume that Conjecture 1.6 holds. Then any compactly supported autoequivalence of $D^b\mathcal{W}(M)$ can be represented by a compactly supported symplectomorphism of M . Moreover, the categorical symplectic mapping class group of M , i.e. the image of the (graded) symplectic mapping class group in $\text{Auteq } D^b\mathcal{W}(M)$, is generated by Lagrangian translations, nodal slide recombinations, and Dehn twists in Lagrangian spheres mirror to line bundles on (-2) curves.*

The theorem holds unconditionally whenever Conjecture 1.6 is known – for instance, for Example 1.3, for which $Z = \emptyset$.

Spherical objects. While our primary objective is not to study spherical objects, our results can be applied to prove some interesting properties.

For instance, Corollary 7.6 shows that whenever there are infinitely many (-2) curves, we can construct a countably infinite family of Lagrangian spheres, none of which is contained in the subcategory of $D^b\mathcal{W}(M)$ split-generated by the others. This should be contrasted with the case of Picard–Lefschetz theory for hypersurface singularities. (Note that while $D^b\mathcal{W}(M)$ is finitely split generated, by non-compact objects, for general reasons, there is no a priori reason to expect the compact Fukaya category, $D^\pi\mathcal{F}(M)$, to be finitely split-generated.)

1.5. Monodromy and moduli spaces of complex structures. We end with some motivation and speculations. Why would one expect autoequivalences of $D(Y)$ to have mirrors realised by symplectomorphisms? For our families, we conjecture that all of these symplectomorphisms are induced by monodromy in the complex moduli space for M .

For the finite list of semi-definite cases, M is the Milnor fibre of a simple elliptic singularity. It compactifies to a del Pezzo surface X , with $M = X \setminus E$, for $E \subset X$ a smooth anticanonical. The moduli space of pairs (X', E') , where X' is deformation equivalent to X and $E' \subset X'$ smooth anticanonical, is well understood. Aspects of this will be revisited in Section 7.3.

For now, focus on the case where D is negative definite. There is now no a priori reason to expect the Stein manifold M to be a variety. To get a precise ‘complex moduli space’ for M , consider the smoothing component of the singularity for which M is a Milnor fibre, with the discriminant locus removed; call this \mathbf{S} . Note that one expects monodromy in a smoothing component to always induce compactly supported symplectomorphisms, whereas for the full complex moduli space in the semi-definite case this won’t always be true.

Conjecture 1.8. *Assume that D is negative definite. There is a natural isomorphism:*

$$\mathbf{S} \simeq \left((Q_{\mathbb{R}} + i\mathcal{C}) \setminus \bigcup_{\alpha \in \Phi, k \in \mathbb{Z}} \{x \mid \langle x, \alpha \rangle = k\} \right) / (Q \rtimes \text{Adm})$$

where $Q = \langle D_1, \dots, D_k \rangle^{\perp} \subset \text{Pic } Y$, for D_1, \dots, D_k the irreducible components of D ; \mathcal{C} is the interior of $\overline{K(Y_{\text{gen}})} \cap Q \subset \text{Pic } Y \otimes \mathbb{R} = H_{dR}^2(Y; \mathbb{R})$, where $K(Y_{\text{gen}})$ is the generic Kaehler cone (i.e. the Kaehler cone for a generic Y' in the complex deformation space of Y); Φ is set of all roots in Q ; and Adm is the subgroup of admissible lattice automorphisms of $\text{Pic } Y$, i.e. the ones preserving the $[D_i]$ and \mathcal{C} .

Remark 1.9. For the purpose of computing $\pi_1(\mathbf{S})$, we could replace \mathcal{C} by the larger cone given by one connected component of the positive square cone; heuristically, this is because we know that the strata of the hyperplane arrangements in these open sets are in bijection.

In the cases $k \leq 5$, Conjecture 1.8 is a theorem of Looijenga [Loo81].

The intersection pairing on $\text{Pic } Y$ has hyperbolic signature, \mathcal{C} is contained in the positive square cone, and Adm acts by isometries, which implies that $Q \rtimes \text{Adm}$ acts properly discontinuously on its domain. This means the quotient is well-behaved.

We expect a representation of $\pi_1(\mathbf{S})$ in $\pi_0 \text{Symp}_c(M)$, and, on the mirror side, in $\text{Auteq}_c D(U)$. Loosely, $Q \rtimes \text{Adm}$ gives the action at the level of homology, with Q corresponding to Lagrangian translations ($\leftrightarrow \otimes \mathcal{O}(\mathcal{L})$), and Adm to the actions of nodal slide recombinations ($\leftrightarrow \text{Aut}(Y, D; \text{pt})$) and Dehn twists ($\leftrightarrow \text{Weyl group}$); and loops in the hyperplane complement should give Torelli elements. We expect an elementary loop around a single hyperplane to correspond to the square of a Dehn twist in a known Lagrangian sphere; on the B side, using the previous notation, these are $T_A \cdot \mathcal{A}$.

The action of $Q \rtimes \text{Adm}$ lifts to $\text{Hom}(K(U), \mathbb{C})$, where $K(U) = \mathbb{Z} \oplus \text{Pic}(U)$ is the K -theory of U . We can consider the space

$$\mathbf{S}' := \left(\text{Hom}(K(U), \mathbb{C})^{\circ} \setminus \bigcup_{\hat{\alpha} \in \hat{\Phi}} \{\hat{\alpha}^{\perp} = 0\} \right) / (Q \rtimes \text{Adm})$$

where $\hat{\Phi}$ is the lift of $\Phi \subset Q$ to $K(U)^*$, and $^{\circ}$ denotes the restriction to the \mathbb{C}^* cover of \mathbf{S} , the moduli space of Conjecture 2. Informally, \mathbf{S}' is the moduli space of complex structures on M equipped with a choice of holomorphic volume form Ω . (This could be made precise

by compactifying the cusp to an Inoue surface, which fixes Ω up to a scalar, cf. [Loo81].) Monodromy about \mathbb{C}^* should correspond to the square of the shift.

Conjecture 1.10. *The universal cover $\tilde{\mathbf{S}}'$ of \mathbf{S}' is a connected component of the space of Bridgeland stability conditions on $D^b \text{Coh}(U)$. The natural map to the intermediate covering space which lies inside $\text{Hom}(K(U), \mathbb{C})$, before quotienting by $Q \rtimes \text{Adm}$, is given by mapping to the central charge.*

We expect that there are two connected components of the space of Bridgeland stability conditions, interchanged by the shift functor, mapping to $Q_{\mathbb{R}} \pm i\mathbb{C}$ via the central charge.

Remark 1.11. The reader may be interested in comparing with work of Engel–Friedman in [EF21, Section 5.1]. They construct monodromy diffeomorphisms of the Milnor fibre M of a smoothing of a cusp singularity associated to a negative definite log CY (Y, D) together with a choice of line bundle $\mathcal{L} \in \mathcal{C}$, describing in particular the action on homology. We expect that M is mirror to (Y, D) and these monodromy diffeomorphisms are the Lagrangian translations mirror to $(\cdot) \otimes \mathcal{L}$, associated to loops bounding a disc passing through the origin in the deformation space of the cusp (rather than an arbitrary closed loop).

Organisation of the paper. Section 2 gathers background material on autoequivalences of the derived categories on the B side. Section 3 contains the input we will use from mirror symmetry, including some refinements of results in [HK]. Section 4 sets up Lagrangian translations, including a proof of Theorem 1.2 part (1) at the K -theoretic level. Section 5 constructs Lagrangian spheres mirror to line bundles on (-2) curves, the second key technical ingredient mentioned above; this also allows us to complete the proof of Theorem 1.2 part (1) (Theorem 5.5). Section 6 constructs nodal slide recombinations and proves Theorem 1.2 part (2), and the independence of choices mentioned above. Relations between symplectomorphisms from different families are established in Section 6.4. Finally, Section 7 contains a range of applications: first, the packaging of the new symplectomorphisms as Theorem 7.1 and discussion of examples in Section 7.1; applications of our constructions to questions about spherical objects in Section 7.2; and it ends with extended discussion of the semi-definite case – further constructions from monodromy (Section 7.3) and connections with work of Collins–Jacob–Lin on hyperkaehler structures (Section 7.4).

Notation and conventions. While they will be introduced as we go along, we record standing notational conventions here for the readers' convenience.

B-side. (Y, D) will denote a log Calabi-Yau pair, assumed to have maximal boundary unless otherwise specified; we use U to denote $Y \setminus D$, k for the number of irreducible components D_1, \dots, D_k of D , and (Y_e, D) for the surface in the deformation class of (Y, D) with the distinguished complex structure, i.e. such that the mixed Hodge structure on $Y_e \setminus D$ is split (see exposition in [HK, Section 2.2]). We say that D is negative definite, semi-definite or

indefinite whenever the intersection form associated to the D_i is. (Abusing terminology, we use ‘semi-definite’ to mean strictly semi-definite.) When D is semi-definite, Y_e is rational elliptic, and we denote by $\varpi : Y_e \rightarrow \mathbb{P}^1$ the minimal rational elliptic fibration ($\varpi^{-1}(\infty) = D$).

A-side. The mirror to (Y_e, D) is a Weinstein manifold M , which is the total space of a Lefschetz fibration $w : M \rightarrow \mathbb{C}$. M is the total space of (the equivalence class of) an almost-toric fibration $\pi : M \rightarrow \mathbb{R}^2$. In the semi-definite case, $M = X \setminus E$, where X is a del Pezzo surface of degree k and $E \subset X$ a smooth anti-canonical divisor (elliptic curve).

Acknowledgements. We are grateful to Roger Casals for correspondence about Remark 6.8; Alessio Corti and Wendelin Lutz for discussions related to [CK16, KSC04]; Abigail Ward for discussions and feedback on a draft of this article; and Ivan Smith for explanations of [SS20a] and for detailed comments on an earlier draft.

PH was partially supported by NSF grant DMS-1901970. AK was partially supported by an award from the Isaac Newton Trust.

2. AUTOEQUIVALENCES OF $D^b \text{Coh}(Y)$ AND $D^b \text{Coh}(U)$

2.1. Weyl group and classes of spherical objects. In the wake of [ST01], a well-studied class of autoequivalences of $D(Y)$ are spherical twists. In particular, given a (-2) curve $C \subset Y$, the sheaves $i_* \mathcal{O}_C(a)$, $a \in \mathbb{Z}$, are spherical objects. Let Z be the union of all (-2) curves in Y . (These are either components of D or disjoint from it.) Any spherical object in $D(Y)$ has support on Z , using the arguments in the proof of [Ueh19, Claim 6.3].

Lemma 2.1. *Suppose that $\tilde{Z} \subset Z$ is either a type A chain of (-2) curves or a cycle of (-2) curves. In the latter case, we further assume that either $\tilde{Z} \neq D$ or $Y = Y_e$. Then the spherical objects with support on \tilde{Z} are the images of*

$$\{i_* \mathcal{O}_C(a) \mid a \in \mathbb{Z}, C \subset \tilde{Z} \text{ a } (-2) \text{ curve}\}$$

under spherical twists in themselves.

Proof. When \tilde{Z} is a type A chain of (-2) curves, this is [IU05, Proposition 1.6 and Lemma 4.1]. When \tilde{Z} is a cycle of (-2) curves, we want to use [Ueh16, Proposition 4.4]. This is stated for the case where \tilde{Z} (Z_0 in the notation in [Ueh16]) is a fibre of a relatively minimal rational elliptic fibration on Y . (Note this would imply that D is semi-definite, cf. start of Section 2.3.1.) However, inspecting the proof of [Ueh16, Proposition 4.4], one can check that relative minimality is not required for that statement: it’s enough to check that there exists an elliptic fibration $f : Y \rightarrow \mathbb{P}^1$ such that \tilde{Z} is a fibre of f . Say \tilde{Z} has components C_1, \dots, C_l .

First, by the adjunction formula, we have that $\omega_{\tilde{Z}} = \omega_Y \otimes \mathcal{O}_Y(\tilde{Z})|_{\tilde{Z}}$. Also, the dualizing sheaf $\omega_{\tilde{Z}}$ is trivial, and $\omega_Y|_{\tilde{Z}} = K_Y|_{\tilde{Z}} = -D|_{\tilde{Z}} = 0$. (Here we use $Y = Y_e$ in the case where $\tilde{Z} = D$.) Thus the normal bundle $\mathcal{O}_Y(\tilde{Z})|_{\tilde{Z}}$ of \tilde{Z} in Y is trivial. It then follows from the exact sequence

$$0 \rightarrow \mathcal{O}_Y \rightarrow \mathcal{O}_Y(\tilde{Z}) \rightarrow \mathcal{O}_Y(\tilde{Z})|_{\tilde{Z}} \rightarrow 0$$

that $\dim H^0(\mathcal{O}_Y(\tilde{Z})) = 2$ (note $H^1(\mathcal{O}_Y) = 0$ because Y is rational). Thus the linear system $|\tilde{Z}|$ gives a rational map to \mathbb{P}^1 .

Next, we claim that $|\tilde{Z}|$ is basepoint free. Write $\tilde{Z} = M + F$ for the moving and fixed parts of the divisor \tilde{Z} . If $F \neq 0$ then $M^2 < 0$ (because $\langle C_1, \dots, C_k \rangle$ is negative semidefinite with kernel $\langle \tilde{Z} \rangle$), contradicting M moving. So $F = 0$ and $\tilde{Z} = M$ is moving. Now $\tilde{Z}^2 = 0$ implies that $|\tilde{Z}|$ is basepoint free. Thus the linear system $|\tilde{Z}|$ gives a morphism to \mathbb{P}^1 , with one fiber equal to \tilde{Z} . (This is essentially the same as the proof of the analogous result for K3 surfaces.) \square

Remark 2.2. In general, there could be infinitely many (-2) curves, and also other finite configurations – e.g. whenever $k = 3$, a nodal configuration with Y-shaped dual graph (mirror to the Milnor fibre of a hypersurface cusp).

When D is indefinite, the only possible configurations of (-2) curves are ADE configurations (and finitely many of them): there's a birational morphism which contracts all the internal (-2) curves to ADE singularities and is an isomorphism elsewhere [GHK15a, Lemma 6.9].

We can completely classify classes of sphericals in $K(D(Y))$. First recall that the set of roots Φ consists of classes in $\text{Pic}(Y)$ obtained by parallel transport from an internal (-2) curve in a deformation equivalent pair (Y', D') . (Alternatively, consider the generic ample cone $\mathcal{A}_{\text{gen}} \subset \text{Pic} Y \otimes \mathbb{R}$; the roots are the (-2) classes δ such that $\langle \delta \rangle^\perp \cap \mathcal{A}_{\text{gen}} \neq \emptyset$, see [Fri].) The Weyl group, denoted by W , is the group generated by reflections $s_\alpha(x) = x + \langle x, \alpha \rangle \alpha$ for $\alpha \in \Phi$, acting on $H^2(Y; \mathbb{Z}) = \text{Pic}(Y)$. The set of *simple roots* Δ consists of classes of (-2) curves $C \subset Y_e \setminus D$. By [GHK15b], W is generated by the reflections in elements of Δ , and $\Phi = W \cdot \Delta$. As the spherical twist in $i_* \mathcal{O}_C(a)$ lifts the action of $s_{[C]}$, any root is the first Chern class of a spherical object in $D(Y)$. The converse also holds:

Lemma 2.3. *Suppose $E \in D(Y)$ is a spherical object with support disjoint from D . Then $c_1(E) \in H^2(Y; \mathbb{Z}) \simeq \text{Pic}(Y)$ is a root.*

Proof. Let E be a spherical object in $D(Y)$ with support disjoint from D . Consider framed deformations of E with framing along D . These are controlled by $\text{Ext}^i(E, E(-D))$, $i = 0, 1, 2$. (Note that, since the support of E is disjoint from D , the framed deformations are identified with ordinary deformations, and $\text{Ext}^i(E, E(-D)) = \text{Ext}^i(E, E)$; however, this is not true for deformations of the associated determinant line bundle considered below.)

Consider the trace map from $\text{Ext}^i(E, E(-D))$ to $H^i(\mathcal{O}_Y(-D))$. This corresponds to the map from framed deformations of E to framed deformations of the line bundle $\det E$. The map

$$\text{Ext}^2(E, E(-D)) \rightarrow H^2(\mathcal{O}_Y(-D))$$

is an isomorphism, that is, the obstruction spaces for deformations of E and framed deformations of $\det E$ are identified. Indeed, recalling that $-D = K_Y$, the map is Serre dual to the map $H^0(\mathcal{O}_Y) \rightarrow \text{Hom}(E, E)$, which is an isomorphism since E is spherical. (This is the usual

argument for smoothness of moduli spaces of stable sheaves on Calabi–Yau surfaces due to Mukai, adapted to the log Calabi–Yau / framed case.)

This implies that if $(\mathcal{Y}, \mathcal{D})/(0 \in S)$ is a deformation of (Y, D) over an analytic germ $(0 \in S)$, then E extends to an object \mathcal{E} in $D^b(\text{Coh } \mathcal{Y})$ iff there is a framed deformation of $\det E$ over S . Equivalently, writing \mathcal{L}_s for the line bundle on \mathcal{Y}_s obtained from $\det E$ by parallel transport (noting that $c_1: \text{Pic } \mathcal{Y}_s \rightarrow H^2(\mathcal{Y}_s, \mathbb{Z})$ is an isomorphism since \mathcal{Y}_s is rational), we have $\mathcal{L}_s|_{\mathcal{D}_s} \simeq \mathcal{O}_{\mathcal{D}_s}$ for all $s \in S$. In the terminology of [GHK15b], after choosing a trivialization of the local system over S with fibers $H^2(\mathcal{Y}_s, \mathbb{Z})$, we require that the period point $\phi_{\mathcal{Y}_s}: H^2(Y, \mathbb{Z}) \rightarrow \mathbb{C}^*$ satisfies $\phi_{\mathcal{Y}_s}(c_1(E)) = 1$ for all $s \in S$. (Moreover, in this case the extension \mathcal{E} is uniquely determined since $\text{Ext}^1(E, E) = 0$.) Write $\alpha = c_1(E) \in H^2(Y, \mathbb{Z})$. Note that $\alpha^2 = -2$ by the Riemann-Roch formula; in particular, $\alpha \in H^2(Y, \mathbb{Z})$ is primitive.

By the local Torelli theorem for log Calabi–Yau surfaces [Loo81], there is a small deformation $(\mathcal{Y}_s, \mathcal{D}_s)$ of (Y, D) such that $\ker \phi_{\mathcal{Y}_s} = \mathbb{Z} \cdot \alpha$. Thus E deforms to a spherical object \mathcal{E}_s on \mathcal{Y}_s . (A deformation of a spherical object is necessarily spherical by upper semicontinuity of coherent cohomology and the topological nature of the Euler characteristic.) Recall that any spherical object in $D(Y)$ has support a union of (-2) curves. Moreover, if $C \subset Y \setminus D$ is an internal curve, then $\phi_Y([C]) = 1$. Finally, the support of \mathcal{E}_s is disjoint from the boundary \mathcal{D}_s of \mathcal{Y}_s for all s (since this is an open condition and it holds for $s = 0$). It follows that $\mathcal{Y}_s \setminus \mathcal{D}_s$ contains a unique (-2) curve C with class $\pm\alpha = \pm c_1(E)$. Thus $c_1(E) \in H^2(Y, \mathbb{Z})$ is a root. \square

Remark 2.4. The K -theory $K(\text{Coh } Y)$ is generated by the image of $\text{Coh}(Y)$ under the Chern character. Explicitly,

$$K(\text{Coh } Y) = \{(r, c_1, \text{ch}_2) \in \mathbb{Z} \oplus H^2(Y, \mathbb{Z}) \oplus \frac{1}{2}\mathbb{Z} \mid \text{ch}_2 \equiv \frac{1}{2}c_1^2 \pmod{\mathbb{Z}}\}.$$

(As Y is rational, the algebraic and topological K -theories for Y agree.) The classes of the spherical objects in $K(\text{Coh } Y_e)$ are given by $(0, c_1, \text{ch}_2)$ where $c_1 \in \Phi$ is a root and $\text{ch}_2 \in \mathbb{Z}$. Above we only considered the first Chern class c_1 , but the same argument shows this stronger result. (Note that the spherical twist acts by the same formula on $K(\text{Coh } Y)$ and $\text{ch}(\mathcal{O}_C(n)) = \text{ch}(\mathcal{O}_C) + (0, 0, n)$ using the basis above.)

2.2. Automorphisms of Y . We collect relevant results from [GHK15b, Fri].

2.2.1. Generalities and Torelli theorem. We consider the following subgroups of the group of biholomorphic automorphisms $\text{Aut}(Y)$: let $\text{Aut}(Y, D)$ be the subgroup preserving D , $\text{Aut}(Y, D; \text{cpt})$, the subgroup preserving each component of D ; and $\text{Aut}(Y, D; \text{pt})$, the subgroup fixing D pointwise. ($\text{Aut}(Y, D; \text{cpt})$ clearly has (at worst) finite index in $\text{Aut}(Y, D)$.)

If D is negative definite, $\text{Aut}(Y) = \text{Aut}(Y, D)$; in the negative semi-definite case, $\text{Aut}(Y, D)$ has finite index in $\text{Aut}(Y)$. In either of these cases, provided D contains no (-1) curves, any automorphism of $U = Y \setminus D$ compactifies to an element of $\text{Aut}(Y)$: this is an straightforward exercise in the semi-definite cases, and follows from uniqueness of the minimal resolution of the

corresponding cusp singularities for the negative definite ones. (Note that this is emphatically not true in the indefinite case, see e.g. [CL09].)

Let Adm denote the subgroup of automorphisms of the lattice $\text{Pic}(Y)$ preserving the classes $[D_i]$ and the generic ample cone in $\text{Pic}(Y)$. (See [GHK15b, Definition 1.7]; this is called $\Gamma(Y, D)$ in [Fri].) Adm is the monodromy group of pairs (Y, D) in the sense of [GHK15b, Theorem 5.15]. We have $W \trianglelefteq \text{Adm}$.

Theorem 2.5. [GHK15b, Theorem 5.1]. *Assume Y has the distinguished complex structure within its deformation class. There is an exact sequence*

$$1 \rightarrow N \rightarrow \text{Aut}(Y, D; \text{cpt}) \rightarrow \text{Adm}/W \rightarrow 1$$

where $N = \text{Hom}(N', \mathbb{G}_m)$, and N' the cokernel of the evaluation map $\text{ev} : \text{Pic}(Y) \rightarrow \mathbb{Z}^k$ given by taking intersection numbers with the D_i . Equivalently, $N' = \pi_1(Y \setminus D)$, by considering the long exact sequence of the pair (Y, U) .

N is the subgroup of $\text{Aut}(Y, D; \text{cpt})$ which acts trivially on $H^2(Y; \mathbb{Z})$. It is finite whenever a toric model for Y involves interior blow ups on components corresponding to (at least) two linearly independent toric rays; in particular, it is finite whenever D is negative semi-definite or negative definite. The action of N on (Y, D) can be described explicitly as follows: choose a toric model $(Y, D) \rightarrow (\bar{Y}, \bar{D})$. Let $T \simeq (\mathbb{C}^*)^2$ be the big torus acting on \bar{Y} . Then N is the subgroup of T fixing the points we blow up to obtain (Y, D) , and the action of N on (\bar{Y}, \bar{D}) lifts to an action of N on (Y, D) .

Lemma 2.6. *Assume that $Y = Y_e$. The exact sequence of Theorem 2.5 splits, and we have*

$$\text{Aut}(Y, D; \text{pt}) = \ker\{\text{Aut}(Y, D; \text{cpt}) \rightarrow \text{Aut } D\} \simeq \text{Adm}/W$$

Proof. From the discussion above, we get $N \subset T \subset \text{Aut}^0(D) (\simeq (\mathbb{C}^*)^k)$. Now consider the restriction map $\text{Aut}(Y, D; \text{cpt}) \rightarrow \text{Aut}^0(D)$. We claim that its image lies in N . This uses the fact that $Y = Y_e$: for any $L \in \text{Pic}(Y)$, $L|_D = \mathcal{O}_D(\sum(L \cdot D_i)p_i)$, for some distinguished points $p_i \in D_i$. Now consider an arbitrary $\varphi \in \text{Aut}(Y, D; \text{cpt})$. It follows from the above that $\varphi|_D^*(L|_D) \simeq \varphi^*(L)|_D \simeq L|_D$. On the other hand, as this holds for an arbitrary L , we see that $\varphi|_D \in N$ by [GHK15b, Proposition 2.6], as required. Thus restriction gives a splitting $\text{Aut}(Y, D; \text{cpt}) \rightarrow N$, and we're done. \square

2.2.2. Negative semi-definite case. Assume that D is negative semi-definite and does not contain a (-1) curve, i.e. D is a cycle of (-2) curves. Complex deformation classes of such pairs (Y, D) are completely understood:

Proposition 2.7. [Fri, Propositions 9.15 and 9.16] *Suppose (Y, D) is a log CY surface, and that D is a cycle of k (-2) curves. Then $k \leq 9$, there is a unique deformation type for $k \neq 8$, and two deformation types for $k = 8$ distinguished by $\pi_1(Y \setminus D)$.*

Remark 2.8. $Y \setminus D$ is mirror to $X \setminus E$ where X is a del Pezzo surface of degree $K_X^2 = k$ and E is a smooth anticanonical divisor [AKO06, HK].

The various flavours of automorphism groups are also known explicitly. First, N is known, see [Fri, Proposition 9.16 (iii)]:

$$\pi_1(Y \setminus D) = \begin{cases} \mathbb{Z}/2 \text{ or } 0 & \text{when } k = 8 \\ \mathbb{Z}/3 & \text{when } k = 9 \\ 0 & \text{otherwise} \end{cases}$$

Second, by [Fri, Corollary 9.20], $\text{Adm} = W$ except when $k = 7$, or $k = 8$ and $\pi_1(Y \setminus D) = 0$. In both of these cases, $\text{Adm}/W \simeq \mathbb{Z}$, see [GHK15b, Example 5.6], [GHK15b, Example 5.3] and [Fri, Lemma 9.18 (iii)].

Thus if $Y = Y_e$, in both of these cases, we get that $\text{Aut}(Y, D; \text{pt}) \simeq \mathbb{Z}$. We can describe these automorphisms explicitly using the Mordell-Weil group of Y . As $Y = Y_e$, it is the total space of a minimal rational elliptic fibration $\varpi : Y \rightarrow \mathbb{P}^1$, with $D = \varpi^{-1}(\infty)$ cycle of (-2) curves. (The existence of such an elliptic fibration is codimension one in the complex moduli space: it exists whenever the period map ϕ satisfies $\phi([D]) = 1$, which is certainly true for Y_e . For background on ϕ , see e.g. [HK, Section 2.2].) The fibration ϖ is unique, and admits a holomorphic section. Recall that the Mordell-Weil group $MW(Y, \mathbb{P}^1)$ is the group of holomorphic sections of ϖ , with identity element the reference section s . This acts by translation on each smooth fibre. As the fibration is relatively minimal, this extends to an automorphism of Y , cf. [BHPVdV04, Proposition III.8.4]. From [Shi90, Theorem 1.3], we have

$$MW(Y, \mathbb{P}^1) \simeq \langle F \rangle^\perp / K$$

where F is the class of a smooth fibre of ϖ , $\langle F \rangle^\perp$ is taken inside $\text{Pic}(Y) = H^2(Y; \mathbb{Z})$ with respect to the standard pairing, and K is the subgroup of $H^2(Y; \mathbb{Z})$ generated by all the irreducible components of fibres. One can calculate that $MW(Y, \mathbb{P}^1) \cong \mathbb{Z}$ when $k = 7$, or $k = 8$ and $\pi_1(Y \setminus D) = 0$, in which cases $\text{Aut}(Y, D; \text{pt}) = k \cdot MW(Y, \mathbb{P}^1)$. In other cases it's a finite group.

2.3. Autoequivalences of $D^b \text{Coh}(Y)$.

2.3.1. D negative definite or indefinite.

Lemma 2.9. [Ueh19] *Assume that D is negative definite or indefinite. Then any autoequivalence of $D(Y)$ has Fourier–Mukai kernel with two-dimensional support.*

Proof. We claim that these cases both fall under (iii) of [Ueh19, Theorem 5.3]: namely that $K_Y \neq 0$, and Y does not admit a minimal elliptic fibration. The second part needs checking. Suppose that Y admits a minimal elliptic fibration. Since Y is rational, there is at most one multiple fiber, say of multiplicity $m \geq 1$, by the Kodaira canonical bundle formula. Then

$D = -K_Y = \frac{1}{m}F$, where F is the class of a fibre. Now $D \cdot F = 0$ so D is contained in a fiber, and $D^2 = 0$ so D is equal to a fiber, and thus D is semi-definite, a contradiction.

□

As before, let Z be the union of all (-2) curves in Y , and let $D_Z(Y)$ be the full triangulated subcategory of $D(Y)$ consisting of objects supported on Z . By [Ueh19, Proposition 6.1], there is a well-defined group homomorphism

$$\text{Auteq } D(Y) \rightarrow \text{Auteq } D_Z(Y)$$

induced by restriction. Let $\text{Auteq}^\dagger D_Z(Y)$ denote the image of this map. (This is $\text{Auteq}_{K\text{-equiv}}^\dagger D_Z(Y)$ in [Ueh19], though in the case at hand all autoequivalences are of so-called K -equivalent-type.)

Theorem 2.10. [Ueh19, Theorems 6.6 and 6.8] *There is a short exact sequence*

$$1 \rightarrow \text{Pic}_Z(Y) \rtimes \text{Aut}_Z(Y) \rightarrow \text{Auteq } D(Y) \rightarrow \text{Auteq}^\dagger D_Z(Y) \rightarrow 1 \quad (2.11)$$

where $\text{Pic}_Z(Y) = \{\mathcal{L} \in \text{Pic}(Y) \mid \mathcal{L}|_Z \simeq \mathcal{O}_Z\}$, acting by tensor product, and $\text{Aut}_Z(Y) = \{\varphi \in \text{Aut}(Y) \mid \varphi|_Z = \text{id}_Z\}$, acting by pushforward.

If Z is a disjoint union of cycles and chains of (-2) curves, and $Y = Y_e$ if D is a cycle of (-2) curves, then

$$\text{Auteq } D(Y) = \langle \text{Br}_Z(Y), \text{Pic}(Y) \rangle \rtimes \text{Aut}(Y) \times \mathbb{Z}[1]$$

by Lemma 2.1, where $\text{Br}_Z = \langle T_{\mathcal{O}_C(a)} \mid C \subset Z \text{ a } (-2) \text{ curve, } a \in \mathbb{Z} \rangle$, and $\mathbb{Z}[1]$ denotes shifts.

2.3.2. D negative semi-definite. Assume that there is a rational elliptic fibration $\varpi : Y \rightarrow \mathbb{P}^1$ as above (e.g. if $Y = Y_e$), and let F denote any fibre of ϖ . Every element of $\text{Auteq } D(Y)$ induces an automorphism of $(K(Y), \chi)$, where χ is the Mukai pairing, and, via restriction, an automorphism of $(K(F), \chi)$: see the proof of [Ueh16, Theorem 3.11] and lemmas leading up to it. (For F we are referring to topological K -theory.) $K(F)$ is identified with $H^0(F, \mathbb{Z}) \oplus H^2(F, \mathbb{Z})$ by taking rank and degree, and, as an element of $GL_2(\mathbb{Z})$, an automorphism preserves χ precisely when it has determinant one. Thus we get a map $\Theta : \text{Auteq } D(Y) \rightarrow SL_2(\mathbb{Z})$.

Theorem 2.12. [Ueh16, Theorem 4.1] *Assume that all the reducible fibres of ϖ are of type I_m (i.e. cycles of (-2) curves). Then there is a short exact sequence*

$$1 \rightarrow \langle \text{Br}_Z, \otimes_{\mathcal{O}_Y}(E) \mid E \cdot F = 0 \rangle \rtimes \text{Aut } Y \times \mathbb{Z}[2] \rightarrow \text{Auteq } D(Y) \xrightarrow{\Theta} SL_2(\mathbb{Z}) \rightarrow 1 \quad (2.13)$$

where $\text{Br}_Z = \langle T_{\mathcal{O}_C(a)} \mid C \subset Y \text{ a } (-2) \text{ curve, } a \in \mathbb{Z} \rangle$, and $\mathbb{Z}[2]$ is even degree shifts.

Note that all (-2) curves are contained in reducible fibres, and so Lemma 2.1 applies. When ϖ has singular fibres of other types, a similar short exact sequence is conjectured in [Ueh16], and known up to the same ambiguity as in the negative definite case: ‘extra’ autoequivalences with Fourier–Mukai kernel with support on Z have not been ruled out.

The degree one shift acts as -1 on K theory, and so is not in the kernel of the map to $SL_2(\mathbb{Z})$. The sequence 2.13 doesn't split; this will be clear from the perspective of moduli space monodromies, see Section 1.5. Explicit lifts of $SL_2(\mathbb{Z})$ elements can be given as follows.

Use the basis $([\mathcal{O}_F], [\mathcal{O}_{pt}])$ for $K(F) \simeq \mathbb{Z}^2$, and let $t = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$ and $s = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ be the usual generators for $SL_2(\mathbb{Z})$. Let E_0 be a holomorphic section of ϖ . Then $\otimes_{\mathcal{O}_Y}(E_0) \in \text{Auteq } D(Y)$ satisfies $\Theta(\otimes_{\mathcal{O}_Y}(E_0)) = t$. Let $\Delta \subset Y \times_{\mathbb{P}^1} Y$ be the relative diagonal, $\pi_i : Y \times_{\mathbb{P}^1} Y \rightarrow Y$ be projection to the i th factor, and let \mathcal{P} be the relative Poincaré bundle associated to E_0 :

$$\mathcal{P} = \mathcal{O}_{Y \times_{\mathbb{P}^1} Y}(\Delta - \pi_1^* E_0 - \pi_2^* E_0).$$

Let $\Phi^{\mathcal{P}}$ be the fibrewise Fourier-Mukai transform with kernel \mathcal{P} ; then $\Theta(\Phi^{\mathcal{P}}) = s$, see e.g. [Bri98].

2.3.3. Detecting the identity.

Proposition 2.14. *Assume that (Y, D) is arbitrary negative definite or indefinite; or semi-definite admitting a rational elliptic fibration (e.g. $Y = Y_e$). Assume that $\psi \in \text{Auteq } D(Y)$ induces the identity on $K(Y)$, and that for each (-2) curve $C \subset Y$, ψ fixes $i_* \mathcal{O}_C(-1)$ and $i_* \mathcal{O}_C$ as objects of $D(Y)$. Then $\psi = \varphi_*$, where $\varphi \in N \leq \text{Aut}(Y, D; \text{cpt})$, up to an even shift if Z is empty. In particular, ψ must be the identity autoequivalence if in addition $\iota^* \circ \psi = \iota^*$, where ι is the inclusion $D \rightarrow Y$, and ι^* the derived pullback $D(Y) \rightarrow \text{Perf}(D)$.*

Proof. Let's first assume that D is either negative definite or indefinite. Let C be an arbitrary (-2) curve. The structure sheaves $i_* \mathcal{O}_p$ of points $p \in C$ are in one-to-one correspondence with cones on morphisms from $i_* \mathcal{O}_C(-1)$ to $i_* \mathcal{O}_C$, so the second assumption implies that ψ permutes these structure sheaves. On the other hand, the proof of [Ueh19, Theorem 6.6] implies for any $q \in Y \setminus Z$, $i_* \mathcal{O}_q$ must be mapped to $i_* \mathcal{O}_{q'}$, some $q' \in Y \setminus Z$. From [Huy06, Corollary 5.23], it follows that ψ is in $\text{Pic}(Y) \times \text{Aut}(Y)$. Now use the assumption on the K -theoretic action: first, considering the image of $[\mathcal{O}_Y] \in K(Y)$, we see that $\psi = \varphi_*$, some $\varphi \in \text{Aut}(Y)$; second, the Torelli theorem (Theorem 2.5) then implies that $\varphi \in N$.

When D is negative semi-definite, as before, ψ must permute structure sheaves of points of Z . By the assumption on K -theory, $\psi \in \ker \Theta$. We want to show that ψ is of K -equivalent type. After composing with an automorphism of \mathbb{P}^1 , ψ restricts to an autoequivalence of $Y \setminus Z$ over $V = \mathbb{P}^1 \setminus \varpi(Z)$ (note here Z is the union of the reducible fibers). Now since $\psi \in \ker(\Theta)$ for any smooth fibre $F \subset Y \setminus Z$ the induced autoequivalence of $D(F)$ is a composition of an automorphism, tensor by a line bundle, and even shift by the description of autoequivalences of $D(F)$ for F an elliptic curve; in particular for any $x \in Y \setminus Z$, $\psi(\mathcal{O}_x) = \mathcal{O}_y[2m]$ for some $y \in Y \setminus Z$ and $m \in \mathbb{Z}$. Finally $m = 0$ if $Z \neq \emptyset$ (since $\psi(\mathcal{O}_x)$ is a sheaf for $x \in Z$ and openness of this property). The argument is then similar to the first case.

The criterion for $\psi = \text{Id}$ is immediate from the characterisation of N in Lemma 2.6. \square

We record relations between different categories of known autoequivalences, for later reference:

Lemma 2.15. *Assume that $\varphi \in \text{Aut}(Y)$, $\mathcal{L} \in \text{Pic}(Y)$, and \mathcal{S} is a spherical object in $D(Y)$. Let $T_{\mathcal{S}}$ denote the spherical twist in \mathcal{S} . Then*

- (1) $\varphi^* \circ T_{\mathcal{S}} = T_{\varphi^*\mathcal{S}} \circ \varphi^*$
- (2) $(_ \otimes \mathcal{L}) \circ T_{\mathcal{S}} = T_{\mathcal{S} \otimes \mathcal{L}} \circ (_ \otimes \mathcal{L})$
- (3) $\varphi^* \circ (_ \otimes \mathcal{L}) = (_ \otimes \varphi^*\mathcal{L}) \circ \varphi^*$
- (4) $T_{\mathcal{O}_C(a-1)} \circ T_{\mathcal{O}_C(a)} = _ \otimes \mathcal{O}_Y(C)$, where C is a (-2) curve and $a \in \mathbb{Z}$.

Proof. These are all immediate apart from the final one, for which the calculation is in [IU05, Lemma 4.15 (i) (2)]. \square

Note also that $\text{Br}_Z \cap \text{Aut}(Y) = \text{Pic}(Y) \cap \text{Aut}(Y) = \{\text{Id}\} \in \text{Auteq}(Y)$, see [IU05, Lemma 4.14].

2.4. ‘Compactly supported’ autoequivalences of $D^b \text{Coh}(U)$. Recall that $D(U)$ is equivalent to the derived wrapped Fukaya category of the mirror, $D^b \mathcal{W}(M)$. On that side, any compactly supported symplectomorphism of M induces an automorphism of $D^b \mathcal{F}^{\rightarrow}(w)$ (the mirror to $D(Y)$), and the identity on the fibre Σ of w , whose Fukaya category $\mathcal{F}(\Sigma)$ is mirror to $\text{Perf}(D)$. This motivates the following definition:

Definition 2.16. We say that $\psi \in \text{Auteq} D(U)$ has compact support if there exists $\tilde{\psi} \in \text{Auteq} D(Y)$ such that $\tilde{\psi}$ restricts to ψ on $D(U)$, i.e. we have a commutative diagram:

$$\begin{array}{ccc} D(Y) & \xrightarrow{\tilde{\psi}} & D(Y) \\ \downarrow i^* & & \downarrow i^* \\ D(U) & \xrightarrow{\psi} & D(U) \end{array}$$

and moreover the following diagram commutes:

$$\begin{array}{ccc} D(Y) & \xrightarrow{\tilde{\psi}} & D(Y) \\ & \searrow i^* & \swarrow i^* \\ & \text{Perf}(D) & \end{array}$$

Let $\text{Auteq}_c D(U)$ denote the subgroup of compactly supported autoequivalences.

Let $Q = \langle D_1, \dots, D_k \rangle^\perp \subset \text{Pic}(Y)$. Assuming $Y = Y_e$, we have that $Q = \{\mathcal{L} \in \text{Pic}(Y) \mid \mathcal{L}|_D \simeq \mathcal{O}_D\}$. Let \bar{Q} be the image of Q in $\text{Pic}(U)$. This is isomorphic to Q unless D is semi-definite, in which case we have a short exact sequence

$$0 \rightarrow \langle D \rangle \rightarrow Q \rightarrow \bar{Q} \rightarrow 0.$$

Choosing (for instance) a section s of $\varpi : Y_e \rightarrow \mathbb{P}^1$ determines a splitting of this sequence:

$$Q \simeq \langle D \rangle \oplus (\langle D_1, \dots, D_k \rangle^\perp \subset \langle D, s \rangle^\perp). \quad (2.17)$$

Conjecture 2.18. *Auteq_c $D(U)$ is generated by \bar{Q} , $\text{Aut}(Y, D; pt)$ and $\text{Br}_{Z'}$, where $Z' = Z \cap U$.*

Whenever Z consists of disjoint chains and cycles of (-2) curves, this follows from Theorems 2.10 and 2.12. In general, the conjecture is known up to rulling out ‘extra’ autoequivalences with support on Z (i.e. restricting trivially to $Y \setminus Z$).

3. MIRROR SYMMETRY FOR LOG CALABI-YAU SURFACES: BACKGROUND

3.1. **Homological mirror symmetry.** Recall the following.

Theorem 3.1. [HK, Theorem 1.1] *Suppose (Y, D) is a log Calabi–Yau surface with maximal boundary, and distinguished complex structure. Then there exists a four-dimensional Weinstein domain M and a Lefschetz fibration $w : M \rightarrow \mathbb{C}$, with fibre Σ , such that:*

- (1) Σ is a k -punctured elliptic curve, where k is the number of irreducible components of D ; there is a quasi-equivalence $D^\pi \mathcal{F}(\Sigma) \simeq \text{Perf}(D)$, due to Lekili–Polishchuk [LP17], where $\mathcal{F}(\Sigma)$ is the Fukaya category of Σ , with objects compact Lagrangian branes. (These are graded categories; in particular, there is a natural choice of line field on Σ .)
- (2) $D^b \mathcal{F}^\rightarrow(w) \simeq D(Y)$, where $\mathcal{F}^\rightarrow(w)$ is the directed Fukaya category of w ;
- (3) $D^b \mathcal{W}(M) \simeq D(Y \setminus D)$, where $\mathcal{W}(M)$ is the wrapped Fukaya category of M .

Remark 3.2. When D is semi-definite, i.e. a cycle of k (-2) curves, M is the Milnor fibre of a rational elliptic singularity, and we have $M = X \setminus E$, where X is a degree k del Pezzo surface, and $E \subset X$ an anti-canonical divisor given by a smooth elliptic curve. [AKO06] proves HMS for these spaces with the A and B sides reversed.

Any log CY pair (Y, D) can be described by the data of a toric model [GHK15a, Proposition 1.3]. We follow [HK, Definition 2.2]. A choice of toric model is given by

$$(\bar{Y}, \bar{D}) \leftarrow (\tilde{Y}, \tilde{D}) \rightarrow (Y, D)$$

where (\bar{Y}, \bar{D}) is a toric pair, the map $(\bar{Y}, \bar{D}) \leftarrow (\tilde{Y}, \tilde{D})$ is given by interior blow ups, and $(\tilde{Y}, \tilde{D}) \rightarrow (Y, D)$ is given by blowing down a sequence of components of \tilde{D} (in other words, one goes from (Y, D) to (\tilde{Y}, \tilde{D}) by performing a sequence of ‘corner’ blow-ups). Varying the blow-up locus for interior blow-ups deforms the complex structure. For the distinguished one, we blow up a single favourite point on each component of \bar{D} , which is determined by the torus action. (See [HK, Section 2.2]. Fixing a torus action, one can take the point $-1 \in \bar{D}_i \setminus \cup_{j \neq i} \bar{D}_j \simeq \mathbb{C}^*$.) When $(\tilde{Y}, D) = (\tilde{Y}_e, D)$, $(\tilde{Y}, \tilde{D}) \rightarrow (\bar{Y}, \bar{D})$ is entirely determined by numerical data:

- the self-intersection numbers of the \bar{D}_i , say n_1, \dots, n_k ;

- the number of interior blow-ups of each component, say m_1, \dots, m_k .

Note also that there's a natural isomorphism $\tilde{Y}_e \setminus \tilde{D} \simeq Y_e \setminus D$; they have the same mirror M , with different choices of superpotential for the two different compactifications (these both give Lefschetz fibrations on M , which are related by a sequence of stabilisations mirror to the sequence of corner blow-ups, see [HK, Proposition 3.11]).

Assume $\tilde{Y} = \tilde{Y}_e$. Given the data of a pair $f : (\tilde{Y}, \tilde{D}) \rightarrow (\bar{Y}, \bar{D})$, we have the following:

- (1) a full exceptional collection for $D(\tilde{Y})$, namely

$$\mathcal{O}_{\Gamma_{km_k}}(\Gamma_{km_k}), \dots, \mathcal{O}_{\Gamma_{k1}}(\Gamma_{k1}), \dots, \mathcal{O}_{\Gamma_{1m_1}}(\Gamma_{1m_1}), \dots, \mathcal{O}_{\Gamma_{11}}(\Gamma_{11}), \mathcal{O}, f^*\mathcal{O}(\bar{D}_1), \dots, f^*\mathcal{O}(\bar{D}_1 + \dots + \bar{D}_{k-1})$$

where Γ_{ij} is the pullback of the j th exceptional curve over \bar{D}_i , for $i = 1, \dots, k$, $j = 1, \dots, m_i$.

- (2) a Lefschetz fibration $w : M \rightarrow \mathbb{C}$ with central fibre a k punctured elliptic curve, and distinguished collection of vanishing cycles

$$\{W_{ij}\}_{i=1, \dots, k, j=1, \dots, m_i}, V_0, \dots, V_{k-1}$$

which correspond under Theorem 3.1 to the full exceptional collection above. We use the notation ϑ_{ij} for the Lefschetz thimble corresponding to W_{ij} .

- (3) a Weinstein deformation equivalence between M and the total space of an explicit almost-toric fibration, below.

Theorem 3.3. [HK, Theorem 1.2] *Let (Y, D) be a log Calabi–Yau surface with maximal boundary and distinguished complex structure. Fix a toric model for (Y, D) , with notation as before. Let v_i be the primitive vector for the ray associated to \bar{D}_i in the fan of \bar{Y} . Then M , the mirror space in Theorem 3.1, is Weinstein deformation equivalent to the total space of an almost-toric fibration with base a two-dimensional integral affine space; smooth fibres Lagrangian two-tori; and a nodal fibre for each of the interior blow-ups on \bar{D}_i , with invariant line in direction v_i . All invariant lines are concurrent. (Note this implies they are cyclically ordered by the indices of the v_i .)*

An explicit identification is constructed in [HK, Section 5.2] for the exact torus mirror to the $(\mathbb{C}^*)^2$ chart ‘inherited’ from $\bar{Y} \setminus \bar{D}$, and then [HK, Section 6.1–2] for the nodal fibres. We often refer to these as ‘Symington almost-toric fibrations’, owing to [Sym03]. One should think of them as SYZ fibrations. The integral affine bases already appear explicitly in [GHK15a]. As a Weinstein handle-body, M is given by attaching two-handles to D^*T^2 ; there is one two-handle for each interior blow-up of a component \bar{D}_i of \bar{D} . The attaching Legendrian in $S(D^*T^2)$ is given as follows: letting N denote the toric lattice, take the line $S^1v_i \hookrightarrow T^2 = N \otimes S^1$ (given

by quotienting $\mathbb{R}v_i \hookrightarrow T^2 = N \otimes \mathbb{R}$, and take its conormal Legendrian lift associated to the orientation induced by v_i . We denote this conormal lift by $S_{v_i^\perp}^1$.

Remark 3.4. Given a toric model for (Y, D) , we can also describe an almost-toric fibration with total space U by starting with the toric fibration $\bar{Y} \rightarrow \mathbb{R}^2$, and adding nodal edge cuts for each of the interior blow ups on \bar{D} , following [Sym03, Section 5.4]. Topologically, the fibrations on U and M as readily dual to each other: in particular, there's an obvious correspondence between nodal fibres, and with respect to the standard bases, the monodromy about one is the transpose of the monodromy about the other, with dual invariant lines. (The reader may recall that as well as being mirror, U and M are diffeomorphic – see e.g. [HK, Remark 1.4].)

3.2. Elementary transformations as almost-toric moves. In [HK, Section 3], we prove that as a Lefschetz fibration, $w : M \rightarrow \mathbb{C}$ is independent of choices. In the present paper, we will later need a strengthening of [HK, Section 6.1.3]: a suitably fine compatibility between changes of toric models for (Y, D) , mutations of the associated mirror Lefschetz fibration $w : M \rightarrow \mathbb{C}$, and changes to the almost-toric fibration on the mirror M (itself also associated to a given choice of toric model for (Y, D)). We establish this in this subsection.

3.2.1. *Change of toric model: B side.*

Proposition 3.5. ([HK, Proposition 3.27]) *Given a pair (Y, D) , any two choices of toric models can be related by a sequence of moves of the following two types:*

- *toric blow ups* [HK, Definition 3.23]. *These are given by performing a toric blow up on (\bar{Y}, \bar{D}) , and the corresponding blow-up on (\tilde{Y}, \tilde{D}) .*
- *elementary transformations* [HK, Definition 3.24]. *Suppose that there are two opposite toric rays in the fan for (\bar{Y}, \bar{D}) , say $v_i = -v_j$, and that $m_i > 0$. The elementary transformation replaces n_i with $n_i - 1$ and n_j with $n_j + 1$, and similarly for m_i and m_j . The other n_l and m_l are unchanged, and (Y, D) itself is unchanged. The toric pair changes, say to $(\bar{Y}^\natural, \bar{D}^\natural)$.*

A toy model for an elementary transformation is between $\mathbb{P}^1 \times \mathbb{P}^1$ (blown up at a point) and the Hirzebruch surface \mathbb{F}_1 (blown up at a point).

This follows from the two-dimensional Sarkisov programme, using the exposition in [KSC04]. It is also closely related to J. Blanc's work on birational automorphisms of $(\mathbb{C}^*)^2$ [Bla13].

3.2.2. *Change of toric model: A side.* Toric blow-ups correspond to stabilisations of the Lefschetz fibration, with a natural compatibility of the respective full exceptional collections, worked out in [HK, Section 3.4.5]. They don't change the almost-toric fibration.

Suppose that $\tilde{Y} = \tilde{Y}_e$, and that we have two different toric models for (\tilde{Y}, \tilde{D}) related by an elementary transformation. The two toric models give two different full exceptional collections

for $D(\tilde{Y})$. We showed that these are related by a sequence of mutations (i.e. Hurwitz moves), and that the two mirror distinguished collections of vanishing cycles are related by the same sequence of mutations [HK, Section 3.4.8].

Assume for convenience that we have $m_j > 1$ and $v_j = -v_k$. (The mutations needed to allow the cyclic reordering of labels are described in [HK, Section 3.4.1].) On the symplectic side, the sequence of Hurwitz moves is encoded in Figure 3.1, and goes as follows. First, mutate W_{jm_j} to the end of the list of vanishing cycles. (Note that by [HK, Proposition 3.15] on the total monodromy of the fibration, this leaves the vanishing cycle itself unchanged.) Then mutate V_{k-1}, \dots, V_j over it to get $V_{k-1}^\natural, \dots, V_j^\natural$. Then mutate W_{jm_j} over V_{j-1}, \dots, V_0 , which gives the vanishing cycle W_k . (We have that $V_l^\natural = V_l$ for $l < j$. This description also ignores trivial Hurwitz moves of meridians over each other.)




FIGURE 3.1. Base of the Lefschetz fibration $w : M \rightarrow \mathbb{C}$, with vanishing paths for the initial distinguished basis (left), and for the one after the elementary transformation (right). \mathbb{W} denotes the collection of all the W_{il} .

We will use the following well-known moves on almost-toric fibrations:

- nodal slide: this deforms the almost-toric fibration by sliding a node along its invariant line (without passing any other nodes that might be on that line), see [Sym03, Section 6.1]. Formally, we have a one-parameter family of exact symplectic manifolds $\{(M_t, \omega_t)\}_{t \in [0,1]}$, which agree outside a compact set, and are symplectically isotopic by Moser.
- cut transfers: for a given node, instead of describing the almost-toric fibration using one invariant half-line, switch to using the other invariant half-line [Via14, Definition 2.1]. This should be thought of as changing the description of an almost-toric fibration rather than the fibration itself; it corresponds to a piecewise linear transformation of the base.

A nodal slide implicitly uses a choice of reference Lagrangian section in a neighbourhood of the nodal fibre. Up to fibrewise Hamiltonian isotopy, there is a unique one locally, and so

this choice is usually suppressed. We will sometimes later need a global choice of Lagrangian section. Note that if we define an almost-toric fibration by starting with $T^*\mathbb{R}^2/\mathbb{Z}^2 \rightarrow \mathbb{R}^2$ and introducing Symington cuts, we get a preferred zero-section, inherited from the zero section in $T^*\mathbb{R}^2$. Unless otherwise specified, we will work with these; they are taken to each other under nodal slides and cut transfers.

Following [HK, Section 6.1.3], elementary transformations are ‘mirror’ to nodal slides combined with cut transfers. We want to make this precise. First, notice that the combinatorics matches up.

Lemma 3.6. *Assume $\tilde{Y} = \tilde{Y}_e$. Suppose $(\tilde{Y}, \tilde{D}) \rightarrow (\bar{Y}, \bar{D})$ and $(\tilde{Y}, \tilde{D}) \rightarrow (\bar{Y}^\natural, \bar{D}^\natural)$ are related by an elementary transformation as above: assume that $v_j = -v_k$ in the fan for \bar{Y} , and that $m_j > 0$, so that we’re replacing n_j with $n_j^\natural = n_j - 1$ and n_k with $n_k^\natural = n_k + 1$, and similarly for the m_l . Start with the mirror almost-toric fibration associated to the first toric model. The one for the second toric model can be obtained by sliding the ‘final’ node (i.e. the one closest to the central fibre) with invariant direction v_j past the central fibre, and then transferring the cut to the $v_k = -v_j$ side.*

Proof. Recall that all elementary transformations are obtained by toric blow ups on one of the ‘model’ ones: the Hirzebruch surfaces \mathbb{F}_a and \mathbb{F}_{a-1} each blown up once for $a \geq 1$ (by convention $\mathbb{F}_0 = \mathbb{P}^1 \times \mathbb{P}^1$) [HK, Proposition 3.37]. This means that it’s enough to check these cases, which can be readily done by hand. \square

The two almost-toric fibrations determine two different Weinstein handbody descriptions for M :

- For the original fibration, start with D^*T , where T is the original (exact) central fibre, i.e. the intersection point of the invariant lines; and, for each nodal fibre with invariant direction v_i , glue a Weinstein two-handle D^*D^2 with attaching Legendrian the half-conormal to $S_{v_i^\perp}^1 \subset T = \mathbb{R}^2/\mathbb{Z}^2$ with coorientation given by v_i . Call these D^*D_{il} , $i = 1, \dots, k$, $l = 1, \dots, m_i$. (For $m_i > 1$, we get an A_{m_i-1} chain of Lagrangian spheres.)
- For the new fibration, start with D^*T' , where T' is the new central fibre, and glue Weinstein two-handles similarly to above, say $D^*D_{il}^\natural$, $i = 1, \dots, k$, $l = 1, \dots, m_i^\natural$.

These are related by a Lagrangian mutation of T over D_{jm_j} . Recall Lagrangian mutations are surgery-type operations modelled on passing from a Clifford to a Chekanov torus; they change the Lagrangian skeleton of the Weinstein domain but not the space itself. These constructions go back to [Aur07], see [PT20, Section 4.4] for a careful general description. (They were first systematically used in [Via14]; we are in the same setting as [STW].) In particular, in our case,

we immediately have that T' is the mutation of T over D_{jm_j} , and under this mutation D_{jm_j} becomes $D_{km_k}^{\natural}$ (this reflects the fact that $v_j = -v_k$), and D_{il} just becomes D_{il}^{\natural} otherwise.

We're now ready to complete our discussion:

Lemma 3.7. *Suppose $(\tilde{Y}, \tilde{D}) \rightarrow (\bar{Y}, \bar{D})$ and $(\tilde{Y}, \tilde{D}) \rightarrow (\bar{Y}^{\natural}, \bar{D}^{\natural})$ are related by an elementary transformation as above. Then the two mirror operations are compatible: under the explicit identification in Theorem 3.3 between M as the total space of the Lefschetz fibration associated to the first toric model, resp. second one, and as the total space of the almost-toric fibration associated to the first toric model, resp. second one, the sequence of Hurwitz moves of Figure 3.1 induces the Lagrangian mutation described above.*

Proof. T is given by iterated Polterovich surgery on the dual collection $V_{k-1}^*, \dots, V_1^*, V_0^*(=V_0)$ [HK, Theorem 5.5]. Moreover, as all elementary transformations are pulled back from model ones between \mathbb{F}_{a-1} and \mathbb{F}_a , inspecting the proof of [HK, Theorem 5.5], we see that the second of the two base cases considering therein always applies for us. Further, we see that the iterated Polterovich surgeries can be factored into gluing V_{k-1}^*, \dots, V_j^* and V_{j-1}^*, \dots, V_0^* , with each giving a cylinder, glued by the product of $W_j \sqcup W_k$ with a small interval. See Figure 3.2. In particular, from this perspective we readily see the handle $D^*D_{jm_j}$, which corresponds to W_{jm_j} under [HK, Proposition 6.3].




FIGURE 3.2. Visualising the Weinstein equivalence between M and the total space of the almost-toric fibration: before the mutation.

Now notice that in the Lefschetz fibration picture, the thimble ϑ_{jm_j} (corresponding to W_{jm_j}) can be Hamiltonian isotoped relative to T to be viewed as the vanishing thimble above the vanishing path for $W_{k,m_k+1} = W_{k,m_k}^{\natural}$, which appears after the Hurwitz moves for the elementary transformation. This isotopy respects the Lefschetz fibration, but not the almost-toric fibration. This is illustrated in Figure 3.3.

The torus T' is given, on the Lefschetz side, by iterated Polterovich surgery on the dual collection $(V_{k-1}^{\natural})^*, \dots, (V_1^{\natural})^*, (V_0^{\natural})^*$, which can be factored as with the description for T . On




FIGURE 3.3. Visualising the Weinstein equivalence between M and the total space of the almost-toric fibration: preparing for the mutation.

the other hand, there is a standard model for a Lagrangian mutation in a Lefschetz fibration: the local model for the mutation between a Clifford and Checkanov torus. This is carefully described in [PT20, Section 4]. By considering that model together with the diagrams for T , T' and $\vartheta_{jm_j} = D_{jm_j}$, we now recognise that T' is given by mutating T over ϑ_{jm_j} . See Figure 3.4. (We've implicitly used the fact that $W_j \cap W_k = \emptyset$ to mutate both of the purple annuli on T in the figure, given by thickenings of the j th and k th meridians, over ϑ_{jm_j} – the second one doesn't change.) \square




FIGURE 3.4. Visualising the Weinstein equivalence between M and the total space of the almost-toric fibration: after mutation. We've isotoped the thimble ϑ_{k,m_k+1} relative to T' a little for legibility.

4. LAGRANGIAN TRANSLATIONS AND TENSORS BY LINE BUNDLES

Suppose M^4 is an exact symplectic manifold, and the total space of an almost-toric fibration $\pi : M \rightarrow B$, as defined in [Sym03], where B is an integral affine manifold homeomorphic to

a disc. We will assume throughout that π only has nodal, i.e. focus-focus, singularities. We also assume that there is a unique nodal singularity in each critical fibre (by sliding along the invariant direction for the singularities, this is true w.l.o.g.). Finally, assume throughout that π is equipped with a reference Lagrangian section, say L_0 . (Recall that this is the case when M is the mirror to a log CY surface with one of the explicit almost-toric fibrations.)

4.1. Spaces of Lagrangian sections. Assume M and π are as above. By the Arnol'd–Liouville theorem on action-angle coordinates [Arn89], any regular fibre F of π has an open neighbourhood that is fibre-preserving symplectomorphic to $(V \times T^2, \omega_0) \rightarrow V$, where $V \subset \mathbb{R}^2$. (Here ω_0 denotes the standard symplectic structure $\sum dp_i \wedge dq_i$ on $\mathbb{R}^2 \times T^2$, where the p_i are the standard coordinates on \mathbb{R}^2 and the q_i the ones on T^2 , normalised to have period 2π .) We'll sometimes refer to patches $V \times T^2$ as above ‘Arnol'd–Liouville charts’. By Duistermaat [Dui80], these can be upgraded to global action-angle coordinates. We briefly recall the features we will use. Consider the map \mathfrak{p}_{L_0} given by:

$$\begin{aligned} \mathfrak{p}_{L_0} : T^*B &\longrightarrow M \\ (q, p) &\longmapsto H_{h_p \cdot \pi}(L_0(q)) \end{aligned}$$

where L_0 is viewed as a map $B \rightarrow M$, h_p is the germ of a smooth function $B(q) \subset B \rightarrow \mathbb{R}$ representing $p \in T_q^*B$, and H_f denotes the time-one Hamiltonian flow of a smooth function f . By construction, \mathfrak{p}_{L_0} intertwines the projection to B . If $q \in B$ is a smooth value of π , then $\mathfrak{p}_{L_0} : T_q^*B \rightarrow \pi^{-1}(q)$ is surjective; if $q \in B$ is a singular value, $\mathfrak{p}_{L_0} : T_q^*B \rightarrow \pi^{-1}(q)$ only misses the critical point. (This uses our assumption that π only has nodal singularities with a single node in each critical fibre.) The period lattice $\Lambda_q^* \subset T_q^*B$ is defined to be $\mathfrak{p}_{L_0}^{-1}(L_0(q))$. This has full rank of q is a smooth value of π , and rank one if q is a critical value. Let B^0 be the complement of the critical values of π in B . Then \mathfrak{p}_{L_0} induces a symplectomorphism $T^*B_0/\Lambda^* \simeq \pi^{-1}(B_0)$. For nodal fibres, we get local symplectomorphisms away from critical points of π .

We want to describe certain spaces of Lagrangian sections of π . Let's start by working away from the critical fibres. As before, let B^0 be the complement of the critical values in B . Let $M^0 = \pi^{-1}(B^0)$ be the complement of the critical fibres, and let $\pi_0 = \pi|_{M^0}$. Let γ denote a fibre of π . By the Leray spectral sequence, there is a short exact sequence

$$0 \rightarrow H^1(R^1(\pi_0)_*\mathbb{Z}) \rightarrow H^2(M^0) \xrightarrow{f_\gamma} \mathbb{Z} \rightarrow 0 \quad (4.1)$$

which automatically splits. On the other hand, by [ABC⁺09, Proposition 6.69], $H^1(R^1(\pi_0)_*\mathbb{Z})$ classifies Lagrangian sections of π_0 up to a fibre-preserving symplectic isotopy, i.e. the addition in T^*B^0 of the graph of a closed one-form on B^0 . In particular, this implies that we have a one-to-one correspondence:

$$\left\{ \begin{array}{l} \text{Lagrangian sections } L \text{ of } \pi_0 \text{ up to} \\ \text{fibre-preserving symplectic isotopy} \end{array} \right\} \longleftrightarrow \left\{ \ker(\cap \gamma : H_2(M^0, \partial M_0) \rightarrow \mathbb{Z}) \right\}$$

$$L \qquad \qquad \qquad \mapsto \qquad \qquad \qquad [L] - [L_0]$$

Let N_i be a small open neighbourhood of the i th singular fibre, and $\partial_i = \partial N_i$ be its boundary; by construction, this is the mapping torus of the transformation $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ acting on γ , for some choice of basis for $H_1(\gamma; \mathbb{Z})$. In particular, $H_1(\partial_i; \mathbb{Z}) \simeq \mathbb{Z} \oplus \mathbb{Z}$, where the first factor is generated by a loop about the critical point in the base, and the second factor is the class of a generator of $H_1(\gamma; \mathbb{Z})$ which intersects the vanishing cycle for this singularity at a single point. (Note that while the class in $H_1(\gamma; \mathbb{Z})$ isn't well-defined, the one in $H_1(\partial_i; \mathbb{Z})$ is.) Projecting to the second factor gives a map $\epsilon_i : H_1(\partial_i; \mathbb{Z}) \rightarrow \mathbb{Z}$.

The Mayer-Vietoris exact sequence and Poincaré–Lefschetz duality imply that

$$H_2(M, \partial M) = \ker \left(\sum \epsilon_i : H_2(M^0, \partial M^0) \rightarrow \bigoplus_i \mathbb{Z} \right). \quad (4.2)$$

We're now ready to fill the critical fibres back in.

Lemma 4.3. *There is a one-to-one correspondence:*

$$\left\{ \begin{array}{l} \text{Lagrangian sections } L \text{ of } \pi \text{ up to} \\ \text{fibre-preserving Hamiltonian isotopy} \end{array} \right\} \longleftrightarrow \left\{ \ker(\frown \gamma : H_2(M, \partial M) \rightarrow \mathbb{Z}) \right\}$$

$$L \qquad \qquad \qquad \mapsto \qquad \qquad [L] - [L_0]$$

As all closed forms on B are exact, fibre-preserving symplectic isotopies are necessarily Hamiltonian.

Remark 4.4. The Leray spectral sequence still implies that there is a short exact sequence

$$0 \rightarrow H^1(R^1(\pi)_* \mathbb{Z}) \rightarrow H^2(M) \xrightarrow{f_\gamma} \mathbb{Z} \rightarrow 0. \quad (4.5)$$

In particular, this characterisation of Lagrangian sections naturally generalises [ABC⁺09, Proposition 6.69].

Proof. of Lemma 4.3. If L is a section of π , then certainly $[L] - [L_0] \in \ker(\frown \gamma : H_2(M, \partial M) \rightarrow \mathbb{Z})$, and two isotopic sections give the same class. Thus the map is well-defined.

Surjectivity: by equation 4.2,

$$\ker(\frown \gamma : H_2(M, \partial M) \rightarrow \mathbb{Z}) \simeq \ker \left((\frown \gamma, \sum \epsilon_i) : H_2(M^0, \partial M^0) \rightarrow \mathbb{Z} \oplus_i \mathbb{Z} \right).$$

Fix an arbitrary class on the right-hand side. There exists a Lagrangian section L of π_0 such that $[L] - [L_0]$ represents that class. It's enough to show that, possibly after a fibre-preserving symplectic isotopy, we can extend L to a section of π .

Let $B_i^0 \subset B^0$ be an open annulus about the i th critical value. From the discussion above, Lagrangian sections of the fibration restricted to B_i^0 , up to fibre-preserving symplectic isotopy, are classified by $\mathbb{Z} = \ker(\frown \gamma : H_2(M_i^0, \partial M_i^0) \rightarrow \mathbb{Z})$; moreover, ϵ_i restricts to an isomorphism $\mathbb{Z} \rightarrow \mathbb{Z}$. It follows that there exist closed forms $\zeta_i \in \Omega^1(B_i^0)$ such that L and L_0 differ by the

graph of ζ_i over B_i^0 . Now pick a closed form $\zeta \in \Omega^1(B^0)$ extending the ζ_i . Let $L' = L - \Gamma(\zeta)$, where $\Gamma(\zeta)$ is the graph of ζ . By construction, L' is symplectic isotopic to L over B^0 via a fibre-preserving map, and it agrees with L_0 on each of the B_i^0 . Now extend L' by L_0 over each of the punctures.

Injectivity: suppose that L and L' are sections of π such that $[L] - [L_0]$ and $[L'] - [L_0]$ agree in $\ker(\cap \gamma : H_2(M, \partial M) \rightarrow \mathbb{Z})$. Then their restrictions to M^0 differ by the graph of a *closed* one-form on B^0 , say ζ . It's enough to show that ζ must also be exact. Let $S_i^1 \subset B_i^0$ be a waist curve of the annulus. Let $V_i : S^1 \times [0, 1] \rightarrow T^*B^0$ be the annulus given by $(\theta, t) \rightarrow t\zeta(\theta)$, for $\theta \in S_i^1$, and let \bar{V}_i be its image in $T^*B^0/\Lambda^* = M$. Now notice that

$$\int_{S_i^1} \zeta = \int_{V_i} \omega = \int_{\bar{V}_i} \omega = \int_{L_0|B_i} \omega - \int_{L|B_i} \omega = 0$$

where $B_i \subset B$ is a disc with boundary S_i^1 , and we are using Stokes' theorem together with the fact that \bar{V}_i , $L_0|B_i$ and $L|B_i$ glue to give a null-homologous two-cycle in M . \square

Remark 4.6. To what extent can we control the behaviour of a section near ∞ ? One can check that for a choice of Liouville form on M suitably adapted to the almost-toric fibration, any Lagrangian section of π can be arranged to be invariant under the Liouville flow outside a compact set, after a fibre-preserving Hamiltonian isotopy. We do not pursue this further here: we expect that a comprehensive treatment of the behaviour of Lagrangian sections near infinity will follow using a notion of monomial admissibility, introduced by Hanlon [Han19], and currently being developed by Hanlon–Ward for mirrors to log CY surfaces [HW].

We haven't yet used exactness of M . However, this is required to prove the next classification.

Proposition 4.7. *Suppose M is an exact symplectic manifold as above, and L_0 a reference Lagrangian section of π . Then we have a one-to-one correspondence:*

$$\left\{ \begin{array}{l} \text{Lag. sections } L \text{ of } \pi \text{ equal to } L_0 \text{ near } \partial M \text{ up to} \\ \text{compactly supported fibre-preserving Ham. isotopy} \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{l} \ker(H_2(M, \partial M) \rightarrow H_1(\partial M)) \\ \simeq H_2(M)/H_2(\partial M) \end{array} \right\}$$

$$L \qquad \mapsto \qquad [L] - [L_0]$$

where the homology groups are taken with integer coefficients.

Proof. The map is clearly well-defined. We have that $H_1(\partial M) \simeq \mathbb{Z} \oplus \mathbb{Z}^2/(A - I)$, where $A \in SL(2, \mathbb{Z})$ is the total monodromy of the fibration; the first factor is given by the base S^1 , and the second by the image of $H_1(\gamma, \mathbb{Z})$. Notice that composing the boundary map with projection to the first factor precisely gives $\cap \gamma : H_2(M, \partial M) \rightarrow \mathbb{Z}$.

Surjectivity: given any class in $\ker(H_2(M, \partial M) \rightarrow H_1(\partial M))$, by Lemma 4.3, there is a Lagrangian section L of π such that $[L] - [L_0]$ is in this class. Further, we have assumed that $[L] - [L_0] \mapsto 0 \in \mathbb{Z}^2/(A - I)$. Now proceed as before: first, notice that again applying the

result for fibrations without singularities [ABC⁺09, Proposition 6.69], the map to $\mathbb{Z}^2/(A - I)$ classifies Lagrangian sections of the restriction of the fibration to a neighbourhood of ∂B , up to addition of the graphs of closed one-forms. Thus L and L_0 differ by the graph of such a one-form, say ζ , near ∂B . Then proceed similarly to the end of the proof of Lemma 4.3: the obstruction to ζ being exact is the symplectic area of an annulus which can be capped off in M by two Lagrangian discs to give a closed cycle. In general, this cycle will no longer be null-homologous. However, by exactness of ω , the obstruction still vanishes. Thus after a fibre-preserving Hamiltonian isotopy (indeed, supported near ∂M), we can assume that L agrees with L_0 near ∂M .

Injectivity: suppose L and L' are such that $[L] - [L'] = 0 \in H_2(M, \partial M)$. By Lemma 4.3, they differ by the graph of an exact one-form; moreover, as they were assumed to agree with L_0 near ∂M , and as ∂B is path-connected, we can take that exact one-form to have compact support. \square

We note the following consequence of the arguments in our proof.

Corollary 4.8. *Suppose L_0, L_1 are Lagrangian sections on π which are smoothly isotopic as sections near ∂B . Then there is a fibre-preserving Hamiltonian isotopy of M , supported near ∂M , such that the image of L_1 agrees with L_0 on a tubular neighbourhood of ∂M .*

4.2. Lagrangian translations.

Proposition 4.9. *Suppose M and π are as above, and L_0, L are Lagrangian sections of π . Let M^s be the smooth locus of π , and let $\mathfrak{p} = \mathfrak{p}_{L_0} : T^*B \rightarrow M^s \subset M$ give global action-angle coordinates, chosen so that \mathfrak{p} takes the zero-section to L_0 . Consider the map*

$$\begin{aligned} \sigma_L : M^s &\rightarrow M^s \\ x &\mapsto \mathfrak{p}(\tilde{x} + \widetilde{L|_{\pi(x)}}) \end{aligned}$$

where \tilde{y} denotes any preimage of $y \in M$ under p . (In words, we are using the linear structure on each fibre to add $L - L_0$.) Then

- (1) σ_L is well defined;
- (2) σ_L extends to a smooth map $M \rightarrow M$, which we also denote by σ_L ;
- (3) σ_L is a symplectomorphism of M ;
- (4) whenever L_0 and L agree near ∂M , σ_L has compact support.

We will refer to σ_L as a *Lagrangian translation*.

Proof. The fact that σ_L is well-defined on M^s is immediate. Locally, L is $\mathfrak{p}(\Gamma(df))$, where f is a smooth real-valued function on a small disc in B ; now notice that σ_L is locally given by the

time-one Hamiltonian flow of $f \circ \pi$. This description makes it clear that σ_L extends over the critical points of π , and that it is a symplectomorphism. The final point is immediate. \square

Remark 4.10. Assume that L_0 and L don't agree near ∂M . Following on Remark 4.6, one could choose the Liouville form on M , and representatives for L_0 and L , so as to get a well-defined action of σ_L on the wrapped Fukaya category $\mathcal{W}(M)$. However, as before, we expect the monomially admissible framework currently being developed in [HW] to be best suited for this.

Let $\pi_0 \text{Symp}_c(M)$ denote the symplectic mapping class group of M . Fix a universal cover $\widetilde{\text{Lag}}$ of the Lagrangian Grassmanian over M . Let $\text{Symp}_c^{\text{gr}}(M)$ be the group of compactly supported, graded symplectomorphisms [Sei00, Section 2.b]. Here we assume both that the underlying diffeomorphism has compact support, and that the bundle map is the identity on $\widetilde{\text{Lag}}$ outside a compact set. (This is a distinguished subgroup of the group of graded symplectomorphisms for which we only require the support of the underlying diffeomorphism to be compact.) By [Sei00, Remark 2.5], there is an exact sequence

$$1 \rightarrow \pi_0 \text{Symp}_c^{\text{gr}}(M) \rightarrow \pi_0 \text{Symp}_c(M) \rightarrow H^1(M, \partial M; \mathbb{Z}) \quad (4.11)$$

where the final map is not a group homomorphism, but measures the difference between the original universal covering $\widetilde{\text{Lag}}$ and its pullback under an element of $\pi_0 \text{Symp}_c(M)$. (Note that in the cases we will consider, M is homotopy equivalent to a CW 2-complex, and in particular $H_3(M; \mathbb{Z}) = 0$.)

We record the following properties of Lagrangian translations, the proofs of which are immediate.

Proposition 4.12. *We get the following:*

- *Lagrangian translations by arbitrary Lagrangian sections give a subgroup of $\pi_0 \text{Symp}(M)$ isomorphic to $\ker(\cap \gamma : H_2(M, \partial M) \rightarrow \mathbb{Z})$.*
- *Lagrangian translations by Lagrangian sections which agree with L_0 near infinity give a subgroup of $\pi_0 \text{Symp}_c(M)$ isomorphic to $H_2(M)/H_2(\partial M)$. Moreover, this lifts to a subgroup of $\pi_0 \text{Symp}_c^{\text{gr}}(M)$.*

The action of σ_L on $H_2(M, \partial M)$ is in either case determined by the fact that σ_L acts on Lagrangian sections by adding $[L] - [L_0]$, and fixes γ and hence the $\mathbb{Z} = \text{Im}(\cap \gamma)$. (Dually, we could use the action on $H^2(M)$, see equation 4.5.) In particular, with respect to the decomposition $(\mathbb{Z}, \ker(\cap \gamma : H_2(M, \partial M) \rightarrow \mathbb{Z}))$, σ_L is given by $\begin{pmatrix} 1 & 0 \\ [L] - [L_0] & \text{Id} \end{pmatrix}$.

We will also want compatibility with nodal slides, which follows from standard Moser-type arguments:

Lemma 4.13. *Suppose that we perform a nodal slide and cut transfer using L_0 as the reference section. Formally, we get almost-toric fibrations $\pi^t : M^t \rightarrow \mathbb{R}^2$, with an isotopy $h^t : M^0 \rightarrow M^t$ preserving the reference Lagrangian sections L_0^t . Suppose $\sigma_L^t \in \pi_0 \text{Symp}_c M^t$ is the Lagrangian translation associated to $[L - L_0^t] \in H_2(M^t)/H_2(\partial M^t)$; then*

$$\sigma_{h_*^t[L]}^t = h^t \circ \sigma_L^0 \circ (h^t)^{-1} \in \pi_0 \text{Symp}_c M^t.$$

Similarly for the non-compactly supported case.

4.3. Mirror symmetry for Lagrangian translations: K -theory. We want to show that Lagrangian translations are mirror to tensors with line bundles. We start by establishing this at the K -theoretic level.

Definition 4.14. Let $\gamma \in H_2(U, \mathbb{Z})$ be the class of a “crossing torus” at a node p of D : identifying the germ $(p \in D \subset Y)$ with $(0 \in (z_1 z_2 = 0) \subset \mathbb{C}_{z_1, z_2}^2)$, γ is the class of the 2-torus $(|z_1| = |z_2| = \epsilon)$ for $0 < \epsilon \ll 1$.

A choice of sign for γ is the same as a choice of generator of $H_1(D, \mathbb{Z})$. Note that γ generates the kernel of the map $i_* : H_2(U, \mathbb{Z}) \rightarrow H_2(Y, \mathbb{Z})$, where $i : U \rightarrow Y$ denotes the inclusion of U in Y .

Lemma 4.15. *Line bundles on U are in one-to-one correspondence, via the first Chern class, with $\ker\left(\int_\gamma : H^2(U) \rightarrow \mathbb{Z}\right) \simeq \ker\left(\frown \gamma : H_2(U, \partial U) \rightarrow \mathbb{Z}\right)$.*

Proof. The long exact sequence for the pair (Y, D) and excision give

$$0 \rightarrow \mathbb{Z}^k \rightarrow H^2(Y) \rightarrow H^2(U) \xrightarrow{\int_\gamma} \mathbb{Z} \rightarrow 0$$

which in turn induces

$$0 \rightarrow \text{Pic}(U) \xrightarrow{c_1} H^2(U) \xrightarrow{\int_\gamma} \mathbb{Z} \rightarrow 0.$$

□

In the semi-definite case, let $F \subset \partial U$ be the class of a torus given by smoothing all the nodes of D . Note that

$$H_2(\partial U) = \begin{cases} \mathbb{Z}\langle\gamma\rangle & \text{in the negative definite case} \\ \mathbb{Z}\langle\gamma\rangle \oplus \mathbb{Z}\langle F\rangle & \text{in the semi-definite case} \end{cases}$$

Now assume $Y = Y_e$. As before, let $Q = \langle D_1, \dots, D_k \rangle^\perp \subset \text{Pic}(Y)$, and let \bar{Q} be the image of Q in $\text{Pic}(U)$.

Lemma 4.16. *The first Chern class gives an isomorphism*

$$\bar{Q} \simeq \ker(H^2(U) \rightarrow H^2(\partial U)) \simeq H_2(U)/H_2(\partial U).$$

Proof. Consider the following commutative diagram

$$\begin{array}{ccccccc}
& & & 0 & & 0 & \\
& & & \downarrow & & \downarrow & \\
0 & \longrightarrow & Q \cap \langle D_1, \dots, D_k \rangle & \longrightarrow & Q & \longrightarrow & \bar{Q} & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \langle D_1, \dots, D_k \rangle & \longrightarrow & H^2(Y) & \longrightarrow & H^2(U) & \xrightarrow{f_\gamma} & \mathbb{Z} \\
& & \text{Id} \downarrow & & (\cdot D_i)_{i=1, \dots, k} \downarrow & & \downarrow & & \text{Id} \downarrow \\
0 & \longrightarrow & H_2(D) & \longrightarrow & H^2(D) \simeq H_2(\nu D, \partial U) \simeq \mathbb{Z}^k & \longrightarrow & H_1(\partial U) & \longrightarrow & H_1(D) \simeq \mathbb{Z}
\end{array}$$

where νD denotes a small neighbourhood of $D \subset Y$ (this deformation-retracts onto D). All rows are exact; the bottom row is part of the long exact sequence for the pair $(\nu D, \partial \nu D) = (\nu D, \partial U)$. Also, all the columns apart from the one involving \bar{Q} are automatically exact; exactness of that final column, and our claim, then follow. \square

Lemma 4.17. $K(U) = K_0(D^b \text{Coh}(U))$ is isomorphic to $\mathbb{Z} \oplus \text{Pic}(U)$ via (rk, c_1) .

Proof. There is an exact sequence

$$K(D) \rightarrow K(Y) \rightarrow K(U) \rightarrow 0$$

using [Har77, Exercise 2.10.6 (b)]. We also have an isomorphism $K(Y) \rightarrow \text{Im}(\text{ch}(Y)) \subset H^{\text{even}}(Y; \mathbb{Q})$, where ch denotes the Chern character; as Y is a surface, this map is given by (rk, c_1, ch_2) . (This follows e.g. from the existence of our exceptional collection for $D(Y)$.)

For each D_i , $K(D_i) \simeq \mathbb{Z}^2$, generated by, for instance, $\mathcal{O}_{D_i}, \mathcal{O}_{p_i}$, where $p_i \in D_i$ can be taken to be any interior point of $D_i \subset D$. Moreover, there is a surjection $\bigoplus_i K(D_i) \rightarrow K(D)$. Applying the Chern character, $\mathcal{O}_{D_i} \mapsto (0, [D_i], -1/2[D_i]^2)$; and all of the \mathcal{O}_{p_i} have the same image, of the form $(0, 0, 1)$. The claim is then clear. \square

We note the following immediate consequence:

Lemma 4.18. Let $\mathcal{L} \in \text{Pic}(U)$. The automorphism $\otimes \mathcal{L} : D^b \text{Coh}(U) \rightarrow D^b \text{Coh}(U)$ induces an automorphism $K(U) \rightarrow K(U)$, given by $\begin{pmatrix} 1 & 0 \\ c_1(\mathcal{L}) & \text{Id} \end{pmatrix}$ on $\mathbb{Z} \oplus \text{Pic}(U)$.

Let's still assume that $Y = Y_e$, and let M be the mirror to $Y \setminus D$, with $\pi : M \rightarrow \mathbb{R}^2$ an almost-toric fibration from Theorem 3.3; note that it only has focus-focus singularities. Let γ_M denote a smooth fibre of π , and on the mirror side, let γ_U denote the class of a crossing torus at a node of D .

Lemma 4.19. *Let $\mathcal{W}(M)$ denote the wrapped Fukaya category of M . Then there is an isomorphism $K_0(D^b\mathcal{W}(M)) \simeq H_2(M, \partial M)$, which takes a Lagrangian brane to \pm its class in relative homology, where the sign depends on the grading of the brane.*

Moreover, we have a commutative diagram of isomorphisms:

$$\begin{array}{ccc} K(U) & \xrightarrow{\text{HMS}} & K_0(D^b\mathcal{W}(M)) \\ \downarrow (c_1, \tau k) & & \downarrow \\ \ker\left(\int_{\gamma_U} : H^2(U) \rightarrow \mathbb{Z}\right) \oplus \mathbb{Z} & \xrightarrow{\iota_*} & \ker(\cap \gamma_M : H_2(M, \partial M) \rightarrow \mathbb{Z}) \oplus \mathbb{Z} \simeq H_2(M, \partial M) \end{array}$$

where the first line is induced by the explicit homological mirror symmetry equivalence between $D^b\text{Coh}U$ and $D^b\mathcal{W}(M)$ in Theorem 3.1, and ι_* is induced by a preferred diffeomorphism $\iota : U \rightarrow M$, which takes γ_U to γ_M , together with Poincaré–Lefschetz duality.

Proof. This follows from assembling previous results. From the description of M in Section 3.1, we see that Lagrangian co-cores generate $H_2(M, \partial M)$; as they also generate the wrapped Fukaya category [CDRGG], the natural map $K_0(D^b\mathcal{W}(M)) \rightarrow H_2(M, \partial M)$ is clearly surjective.

On the other hand, by using the proof of Theorem 3.1 (3) together with descriptions of M given by attaching handles to D^*T^2 along Legendrian (also recalled in Section 3.1), we get a diffeomorphism $\iota : U \rightarrow M$ such that the diagram in our statement above commutes. (This is checked inductively, starting with a toric pair and performing interior blow-ups / handle attachments. See also [HK, Remark 1.4].) As the two horizontal maps and the left-hand vertical one are isomorphisms, the right-hand vertical one must be too. \square

The following corollary is then immediate

Corollary 4.20. *Suppose $(Y, D) = (Y_e, D)$ is a log Calabi–Yau surface with distinguished complex structure, with a choice of toric model, and $w : M \rightarrow \mathbb{C}$ its mirror. Set $U = Y \setminus D$, and let $\pi : M \rightarrow B$ be the almost-toric fibration on M associated to that model, with reference Lagrangian section L_0 .*

(1) *The preferred diffeomorphism $\iota : U \rightarrow M$ above induces identifications:*

$$\begin{array}{ccc} \left\{ \begin{array}{l} \text{Lagrangian sections } L \text{ of } \pi \text{ up to} \\ \text{fibre-preserving Hamiltonian isotopy} \end{array} \right\} & \longleftrightarrow & \left\{ \ker(\cap \gamma_M : H_2(M, \partial M) \rightarrow \mathbb{Z}) \right\} \\ & & \updownarrow \\ \text{Pic}(U) & \longleftrightarrow & \left\{ \ker\left(\int_{\gamma_U} : H^2(U) \rightarrow \mathbb{Z}\right) \right\} \end{array}$$

$$\left\{ \begin{array}{l} \text{Lagrangian sections } L \text{ of } \pi \text{ equal to } L_0 \text{ near } \partial M \\ \text{up to fibre-preserving Hamiltonian isotopy} \end{array} \right\} \longleftrightarrow H_2(M)/H_2(\partial M)$$

$$\begin{array}{ccc} & & \updownarrow \\ & & H_2(U)/H_2(\partial U) \\ \text{Line bundles in } \bar{Q} & \longleftrightarrow & \end{array}$$

- (2) Suppose that a section L corresponds to a line bundle \mathcal{L} in \bar{Q} . As an element of $\pi_0 \text{Symp}_c^{\text{gr}}(M)$, σ_L acts on $\mathcal{W}(M)$, and the induced action on $K_0 D^b \mathcal{W}(M)$ naturally agrees with the homological action $[\sigma_L]_*$ on $H_2(M, \partial M; \mathbb{Z})$. Moreover, under the identification given by homological mirror symmetry (as in Lemma 4.19), the action of $\otimes \mathcal{L}$ on $K(U)$ agrees with the action of σ_L on $K_0 D^b \mathcal{W}(M)$.

Note that even in the non-compact case we can always make sense of the K -theoretic action up to a shift (depending on the choice of lift of σ_L to a graded Lagrangian), and that the shift can be chosen to get it to agree with the action of $\otimes \mathcal{L}$.

5. LAGRANGIAN SPHERES MIRROR TO (-2) CURVES

5.1. Mirrors to line bundles on (-2) curves: construction.

Proposition 5.1. *Let (Y, D) be a log CY surface, and let $C \subset Y \setminus D$ be a (-2) curve. Then there exists a toric model for (Y, D) that contracts C : in other words, C is the strict transform of the exceptional divisor for an interior blow-up.*

Proof. This follows from the minimal model program for surfaces, and was noted by Corti–Filip–Petracci as [CFP, Lemma 3.7]. \square

This can be combined with results from Section 3.1 to give the following.

Proposition 5.2. *Assume that (Y, D) is a log CY surface with $Y = Y_e$, and M the mirror to $U = Y \setminus D$. Consider the explicit homological mirror symmetry equivalence between $D^b \text{Coh } U$ and $D^b \mathcal{W}(M)$ given by Theorem 3.1 (3). Suppose $C \subset U$ is a (-2) curve. Then*

- (1) *There exists an embedded Lagrangian sphere $s_C \subset M$ such that s_C is mirror to $i_* \mathcal{O}_C(-1)$. This can be explicitly described as a matching cycle for the Lefschetz fibration $w : M \rightarrow \mathbb{C}$, and, for a suitable choice of almost-toric fibration on M , s_C lies above a segment joining two nodal critical points with the same invariant direction.*
- (2) *For any $E \subset Y$ such that $E \cdot C = 1$, let σ_E be the Lagrangian translation associated to $\otimes \mathcal{O}(E)$. Then for any $a \in \mathbb{Z}$, $\sigma_E^a s_C$ is mirror to $i_* \mathcal{O}_C(a - 1)$.*

Proof. First, notice that the independence of choice of toric model of the mirror, as established in [HK, Section 3.4], means that we are allowed to work with a different toric model for each (-2) curve.

Assume now that our (-2) curve $C \subset U$ arises as part of a toric model $f : (\tilde{Y}, \tilde{D}) \rightarrow (\bar{Y}, \bar{D})$, where (\bar{Y}, \bar{D}) is toric, and (\tilde{Y}, \tilde{D}) is obtained from (Y, D) by a sequence of corner blow-ups (so $U = Y \setminus D \simeq \tilde{Y} \setminus \tilde{D}$). Consider the usual exceptional collection on \tilde{Y} , namely

$$\begin{aligned} \mathcal{O}_{\Gamma_{km_k}}(\Gamma_{km_k}), \dots, \mathcal{O}_{\Gamma_{k1}}(\Gamma_{k1}), \dots, \mathcal{O}_{\Gamma_{1m_1}}(\Gamma_{1m_1}), \dots, \mathcal{O}_{\Gamma_{11}}(\Gamma_{11}), \mathcal{O}, f^*\mathcal{O}(\bar{D}_1), \\ \dots, f^*\mathcal{O}(\bar{D}_1 + \dots + \bar{D}_{k-1}) \end{aligned} \quad (5.3)$$

where Γ_{ij} is the pullback of the j th exceptional curve over \bar{D}_i , for $i = 1, \dots, k$, $j = 1, \dots, m$.

Let's first prove the claim about $i_*\mathcal{O}_C(-1)$ in (1). Assume that C is the strict transform of the j th exceptional curve over \bar{D}_i , for some fixed i, j . Then

$$i_*\mathcal{O}_C(-1) \simeq \{e : \mathcal{O}_{\Gamma_{i(j+1)}}(\Gamma_{i(j+1)}) \rightarrow \mathcal{O}_{\Gamma_{ij}}(\Gamma_{ij})\}$$

where e is a non-trivial constant in $\text{Hom}(\mathcal{O}_{\Gamma_{i(j+1)}}(\Gamma_{i(j+1)}), \mathcal{O}_{\Gamma_{ij}}(\Gamma_{ij})) \simeq \mathbb{C}$.

On the mirror side, $\mathcal{O}_{\Gamma_{i(j+1)}}(\Gamma_{i(j+1)})$ and $\mathcal{O}_{\Gamma_{ij}}(\Gamma_{ij})$ correspond to consecutive thimbles $\vartheta_{i,j+1}$ and $\vartheta_{i,j}$ with the same vanishing cycle, namely the i th meridian (see e.g. [HK, Section 3.3.2]). The morphism e corresponds to a non-trivial degree zero morphism in $H^1(S^1) \simeq HF(\vartheta_{i,j+1}, \vartheta_{i,j})$, where we are working in the directed Fukaya category of the superpotential. The cone over this element is the matching cycle given by gluing $\vartheta_{i,j+1}$ and $\vartheta_{i,j}$ together, by [Sei08, Proposition 18.21]. Let this be the Lagrangian sphere s_C .

The assertion about matching cycles for the Lefschetz fibration is satisfied by construction. For the one about almost-toric fibrations, consider the almost-toric fibration associated to the same choice of toric model. The fibration is described in Theorem 3.3; inspecting the construction of the symplectomorphism between it and M in [HK, Section 6], we see that s_C is a Lagrangian sphere lying above the j th segment between the critical values with invariant direction the ray for \bar{D}_i . (See [Sym03, Section 7.1] for a more general discussion of such Lagrangians.)

We now want to prove the claim about $i_*\mathcal{O}_C(a-1)$ in (2). Set $f = f_2 \circ f_1$, where f_1 is given by starting (\bar{Y}, \bar{D}) and blowing up the i th component of the divisor j times, to get the log CY pair (Y^\dagger, D^\dagger) , say, and f_2 is given by making all the remaining blow-ups. Now notice there is an isomorphism

$$i_*\mathcal{O}_C(a-1) \simeq \{e : \mathcal{O}_{\Gamma_{i(j+1)}}(\Gamma_{i(j+1)}) \rightarrow \mathcal{O}_{\Gamma_{ij}}(\Gamma_{ij}) \otimes f_2^*(\mathcal{O}(D^\dagger))^{\otimes a}\}.$$

On the other hand, $\mathcal{O}_{\Gamma_{ij}}(\Gamma_{ij}) \otimes f_2^*(\mathcal{O}(D^\dagger))$ is given by mutating $\mathcal{O}_{\Gamma_{ij}}(\Gamma_{ij})$ over $f^*\mathcal{O}(\bar{D}_1), \dots, f^*\mathcal{O}(\bar{D}_1 + \dots, \bar{D}_{k-1})$: this follows from applying [BS10, Corollary 2.10] to the exceptional collection for (Y^\dagger, D^\dagger) , and pulling everything back under f_2 . (We can ignore the $\Gamma_{i'j'}$ for $i' \neq i$.) This is the same as mutating it over the dual exceptional collection $f^*(\mathcal{O}(\bar{D}_1 + \dots, \bar{D}_{k-1})^*), \dots, f^*(\mathcal{O}(\bar{D}_1)^*), \mathcal{O}$. Let $\varsigma_{k-1}, \dots, \varsigma_0$ be the mirror thimbles. These can be glued together by iterative Polterovich surgery to get the exact Lagrangian torus in the copy of $(\mathbb{C}^*)^2$ mirror to $\bar{Y} \setminus \bar{D}$ (see [HK, Theorem 5.5]). Let $\vartheta_{i,j}$ be as above.

Now inspect the proof of [HK, Proposition 6.3]: notice that the vanishing cycle $W_i = \partial(\vartheta_{i,j})$ intersects the vanishing cycle $\partial\varsigma_{k-1}$ transversally at a single point; moreover, the Polterovich surgery $\partial(\vartheta_{i,j})\#\partial\varsigma_{k-1}$ intersects $\partial\varsigma_{k-2}$ transversally at a single point, and the same remains true as one iterates. On the other hand, in the case where two thimbles intersect transversally at a single point, mutating one over the other is the same as performing Polterovich surgery (c.f. [Sei99] and [Sei08, Section 18]); this also implies that the result of this iterative surgery has boundary a vanishing cycle which is still a copy of the i th meridian, by applying the total monodromy of the part of the fibration that is mirror to (\bar{Y}, \bar{D}) , see [HK, Proposition 3.15]. It follows that $\mathcal{O}_{\Gamma_{ij}}(\Gamma_{ij}) \otimes f^*(\mathcal{O}(D^\dagger))$ is mirror to a sphere given by performing an iterated Polterovich surgery on $\vartheta_{i,j}, \varsigma_{k-1}, \dots, \varsigma_0$, and then capping off the boundary of the resulting Lagrangian with $\vartheta_{i,j+1}$; topologically, this is clearly a Lagrangian sphere; call it s'_C . To relate s'_C to $\sigma_E(s_C)$, recall that we also know that iterated Polterovich surgery on the ς_l gives an exact Lagrangian torus T which can be taken to be a fibre of the Symington almost-toric fibration; and $\vartheta_{i,j}$ and $\vartheta_{i,j+1}$ are locally given by half-conormals to $S^1_{v_i^\perp}$ in $D_\epsilon^*T^2$, where $v_i \in \mathbb{R}^2$ is the ray associated to \bar{D}_i in the fan for (\bar{Y}, \bar{D}) , and $S^1_{v_i^\perp}$ the image of $\mathbb{R}v_i^\perp$ in $T^2 = \mathbb{R}^2/\mathbb{Z}^2$ (see Section 3.1). In particular, we see that s_C can be arranged to cleanly intersect T^2 along $S^1_{v_i^\perp}$, and that s'_C is the result of an S^1 Lagrangian surgery between them (with suitable sign). Now notice that $\sigma_E(s_C)$ can be described in precisely the same way.

For $\sigma_E^{-1}(s_C)$, one can proceed similarly, taking the Polterovich surgery (with the other ordering choice) of $\vartheta_{i,j}$ with ς_0 , then ς_1 , etc, which will correspond to twisting $\mathcal{O}_{\Gamma_{ij}}(\Gamma_{ij})$ with $\mathcal{O}(-D)$, and taking the S^1 Lagrangian surgery between s_C and T^2 with the opposite sign to before. More generally, notice that one can iterate this process by using disjoint parallel copies of T^2 in $D_\epsilon^*(T^2)$. (At most one of them can be exact, but this doesn't matter after surgery to get a Lagrangian sphere.) \square

Remark 5.4. While we won't need this in this paper, we expect that up to Hamiltonian isotopy, the Lagrangian sphere s_C constructed above is independent of the choice of toric model contracting C . This would follow e.g. from a strengthening of Proposition 3.5: we expect that the sequence of moves between any two toric models contracting C can be chosen so that C is contracted at each step. (For these purposes we consider any power of a fixed elementary transformation to be a single step.) Using the discussion of mirror moves in Section 3.2.2, we would then get that under the corresponding sequence of nodal slides and cuts, s_C would remain a matching cycle between two nodal critical points at all time.

5.2. Full mirror symmetry for Lagrangian translations.

Theorem 5.5. *Suppose $(Y, D) = (Y_e, D)$ is a log Calabi-Yau surface with distinguished complex structure, and $w : M \rightarrow \mathbb{C}$ its mirror. Set $U = Y \setminus D$, let $\pi : M \rightarrow B$ be an almost-toric fibration on M , and L_0 a reference Lagrangian section of π . Assume that L is a Lagrangian section of π which agrees with L_0 near ∂M , and $\mathcal{L} \in \bar{Q} \subset \text{Pic } U$ the corresponding line bundle, as in*

Corollary 4.20. *Then under the homological mirror symmetry equivalence $D(U) \simeq D^b\mathcal{W}(M)$ of Theorem 3.1 (3), $\otimes\mathcal{L}$ corresponds to σ_L .*

Proof. As it is compactly supported, $\sigma_L \in \pi_0 \text{Symp}_c^{\text{gr}}(M)$ gives an autoequivalence of $D^b\mathcal{F}^\rightarrow(w)$ (using e.g. the characterisation of $D^b\mathcal{F}^\rightarrow(w)$ in [GPS20]: we can define the category using exact Lagrangians with prescribed behaviour in a neighbourhood of the boundary but no conditions inside of this). Moreover, this restricts to the identity on $D^\pi\mathcal{F}(\Sigma)$, where Σ is a fibre of w near infinity. Under mirror symmetry, we get an autoequivalence of $D(Y)$ which restricts to the identity on $\text{Perf } D$, say $\check{\sigma}_L$. Consider $\check{\sigma}_L^{-1} \circ (_ \otimes \mathcal{L}) \in \text{Auteq } D(Y)$. First, Corollary 4.20 implies that it acts as the identity on K -theory of $D(U)$; also, Proposition 5.2, together with Lemma 4.13, tells us that it acts as the identity on $\mathcal{O}_C(a)$ for any (-2) curve C in U and any $a \in \mathbb{Z}$. The proof of Proposition 2.14, together with the classification result for chains of (-2) curves (Lemma 2.1, then constrains $\check{\sigma}_L^{-1} \circ (_ \otimes \mathcal{L})$ as an element of $\text{Auteq } D(Y)$: it could only be a product of spherical twists in objects $\mathcal{O}_{C'}(a)$ for $C' \subset D$ a (-2) curve. In particular, it restricts to the identity in $\text{Auteq } D(U)$. \square

Remark 5.6. If D is indefinite or negative definite, we have $Q = \bar{Q}$, and it follows from the proof above that $\otimes\mathcal{L}$ corresponds to σ_L under the equivalence $D(Y) \simeq D^b\mathcal{F}^\rightarrow(w)$. In the negative semi-definite case, there is an overall ambiguity of $\otimes\mathcal{O}(D)$, i.e. the action of the Serre functor. The mirror to this is well-understood, going back to [Kon]: it's action of the total monodromy of the Lefschetz fibration on the directed Fukaya category, cf. [Sei17] and follow-up papers. In the toric case, this was also carefully studied by Hanlon [Han19].

Remark 5.7. In the non-compact case, if we knew that σ_L acted both on $\mathcal{W}(M)$ and $D^b\mathcal{F}^\rightarrow(w)$, then we would be able to apply Proposition 2.14 to show that σ_L is mirror to tensoring with \mathcal{L} . (This would make use of case (2) of the proposition, by considering the action of σ_L on the central torus.) This crucially relies on having an action on $D^b\mathcal{F}^\rightarrow(w)$, as the Prop. 2.14 applies to autoequivalences of $D(Y)$. In order to do this, we need a condition on the behaviour of L near infinity that is more rigid than simply being Liouville adapted (reflecting the difference between $\text{Pic } Y$ and $\text{Pic } U$), and we again refer the reader to the forthcoming work of Hanlon–Ward on monomial admissibility [HW].

6. NODAL SLIDE RECOMBINATIONS

Assume throughout this section that $(Y, D) = (Y_e, D)$.

6.1. Construction and mirror theorem. Fix a toric model $(\bar{Y}, \bar{D}) \leftarrow (\tilde{Y}, \tilde{D}) \rightarrow (Y, D)$. Suppose $\varphi \in \text{Aut}(Y, D; \text{pt})$ is an automorphism of Y fixing D pointwise; this induces an automorphism of \tilde{Y} fixing \tilde{D} pointwise, which we also denote φ . Let $f : (\tilde{Y}, \tilde{D}) \rightarrow (\bar{Y}, \bar{D})$ denote the blow-down map. Now consider the map $f \circ \varphi : (\tilde{Y}, \tilde{D}) \rightarrow (\bar{Y}, \bar{D})$. This also gives a toric model for (\tilde{Y}, \tilde{D}) (and indeed, (Y, D)). By Proposition 3.5, one can go between these via blow-ups and elementary transformations, determined by a factorisation of φ into elementary

birational transformations of (\tilde{Y}, \tilde{D}) . After performing toric blow ups to (\tilde{Y}, \tilde{D}) , we can assume wlog that we're only making elementary transformations, say $\mathcal{E}_1, \dots, \mathcal{E}_r$. Notice that combinatorially, on the mirror side, we 'get back' to the Lefschetz fibration, and almost toric fibration, that we started with: the n_i and m_i are the same at the start and at the end.

Theorem 6.1. *Fix $\varphi \in \text{Aut}(Y, D; pt)$, and a factorisation of it into elementary transformations, say $\mathcal{E}_1, \dots, \mathcal{E}_r$, as above. Then there exists a compactly supported symplectomorphism $\tilde{\varphi} \in \pi_0 \text{Symp}_c^{\text{gr}}(M)$ such that:*

- (1) *With respect to the Lefschetz fibration $w : M \rightarrow \mathbb{C}$, $\tilde{\varphi}$ is induced by a compactly supported symplectomorphism of the base, relative to the singular values, determined by the full sequence of Hurwitz moves for $\mathcal{E}_1, \dots, \mathcal{E}_r$;*
- (2) *With respect to the almost-toric fibration on M associated to our choice of toric model, $\tilde{\varphi}$ is induced by the sequence of nodal slides and cut transfers determined by $\mathcal{E}_1, \dots, \mathcal{E}_r$ (using the standard reference Lagrangian section L_0 , itself fixed).*
- (3) *Under the homological mirror symmetry identification (Theorem 3.1 (2) and (3)), the action of φ_* on the category $D(Y)$, respectively $D(U)$, is mirror to the action of $\tilde{\varphi}$ on the category $D^b\mathcal{F}^\rightarrow(w)$, respectively $D^b\mathcal{W}(M)$.*

Proof. Let's start with (1). Consider the explicit Lefschetz fibration associated to our choice of toric model. Using the same notation as before, it has a distinguished collection of vanishing cycles $\{W_{ij}\}_{i=1, \dots, k, j=1, \dots, m_i}$, V_0, \dots, V_{k-1} , with vanishing paths, say, $\{\gamma_{ij}\}, \varrho_0, \dots, \varrho_{k-1}$. Now perform the Hurwitz moves for $\mathcal{E}_1, \dots, \mathcal{E}_r$. (This follows Figure 3.1.) At then end, by our observation that the n_i and m_i are the same as at the start, the distinguished collection that we get has, again, ordered vanishing cycles $\{W_{ij}\}_{i=1, \dots, k, j=1, \dots, m_i}$, V_0, \dots, V_k . The vanishing paths, however, will have changed (indeed, the ordering of the critical points that they induce should in general also have changed). Call the new paths $\{\gamma'_{ij}\}, \varrho'_0, \dots, \varrho'_{k-1}$.

Up to compactly supported isotopy relative to the critical points, there exists a unique compactly supported orientation-preserving diffeomorphism of the base taking the γ_{ij} to γ'_{ij} , the ϱ_l to the ϱ'_l , and preserving a straight half-line going to $-\infty$ out of the central fibre. (Before requiring the latter, the map is only defined by to the ambiguity of rotating by $2k\pi$ in a large radius annulus.) Call this map $f_{\tilde{\varphi}}$. Similarly to the construction in [Kea, Proposition 2.5], this induces a compactly supported symplectomorphism $\tilde{\varphi}$ of the total space M , uniquely determined up to Hamiltonian isotopy, such that

- on a large compact set K with $f(K) \supset \text{Supp } f_{\tilde{\varphi}}$, it intertwines w , and lifts $f_{\tilde{\varphi}}$;
- in a neighbourhood of the vertical boundary of the Lefschetz fibration above $f(K_1)$, it interpolates back between $f_{\tilde{\varphi}}$ and the identity (recall that after forgetting the marked points, $f_{\tilde{\varphi}}$ is isotopic to the identity)

- it's the identity on fibres above points outside of $f(K)$.

This completes (1). (The reader may also be interested in the set-up at the end of [Tor, Section 3] in a related setting.)

From the definition of $\check{\varphi}$, it maps the Lagrangian branes supported on the thimbles above $\{\gamma_{ij}\}, \varrho_0, \dots, \varrho_{k-1}$ to the ones above $\{\gamma'_{ij}\}, \varrho'_0, \dots, \varrho'_{k-1}$. On the other hand, under the HMS theorem, the former are mirror to

$$\mathcal{O}_{\Gamma_{km_k}}(\Gamma_{km_k}), \dots, \mathcal{O}_{\Gamma_{k1}}(\Gamma_{k1}), \dots, \mathcal{O}_{\Gamma_{1m_1}}(\Gamma_{1m_1}), \dots, \mathcal{O}_{\Gamma_{11}}(\Gamma_{11}), \mathcal{O}, f^*\mathcal{O}(\bar{D}_1), \\ \dots, f^*\mathcal{O}(\bar{D}_1 + \dots + \bar{D}_{k-1})$$

where $f : (\tilde{Y}, \tilde{D}) \rightarrow (Y, D)$ is part of the toric model datum. On the other hand, using the fact that the elementary transformations $\mathcal{E}_1, \dots, \mathcal{E}_r$ are factorising φ , the branes supported on the thimbles above $\{\gamma'_{ij}\}, \varrho'_0, \dots, \varrho'_{k-1}$ are mirror to

$$\mathcal{O}_{\varphi(\Gamma_{km_k})}(\varphi(\Gamma_{km_k})), \dots, \mathcal{O}_{\varphi(\Gamma_{k1})}(\varphi(\Gamma_{k1})), \dots, \mathcal{O}_{\varphi(\Gamma_{1m_1})}(\varphi(\Gamma_{1m_1})), \dots, \mathcal{O}_{\varphi(\Gamma_{11})}(\varphi(\Gamma_{11})), \\ \mathcal{O}, f^*\mathcal{O}(\bar{D}_1), \dots, f^*\mathcal{O}(\bar{D}_1 + \dots + \bar{D}_{k-1})$$

where we are using the fact that the \bar{D}_i are fixed by φ . This completes (3).

For (2), consider the sequence of nodal slides and cut transfers induced by $\mathcal{E}_1, \dots, \mathcal{E}_r$. We claim that as the n_i and m_i at the end are the same as the ones we started with, the sequence of almost-toric moves naturally induces a compactly supported symplectomorphism of M . Formally, this can be defined as follows. Set $M = M_0$. The first nodal slide gives a one-parameter family of symplectic manifolds M_t , $t \in [0, 1]$, naturally identified outside a compact set, with a smoothly varying family of symplectomorphisms $\check{\varphi}_t : M_0 \rightarrow M_t$, all compactly supported (this makes sense because of the natural identification), and such that $\check{\varphi}_0 = \text{Id}$. The first cut transfer simply takes M_1 to itself (no deformations). The second nodal slide extends our families to M_t and $\check{\varphi}_t$ with $t \in [1, 2]$. Iterating, we get smoothly varying families M_t and $\check{\varphi}_t$ for $t \in [0, r]$ (wlog we reparametrise near integer points of $[0, r]$ to get smoothness everywhere). Now, after all r steps, the n_i and m_i are the same as we started with; thus M_r is naturally identified, as a symplectic manifold, with $M = M_0$; and $\check{\varphi}_r$ is a compactly supported symplectomorphism of M .

We claim that under our identification of M as the total space of the Lefschetz fibration w and as the total space of the (initial) almost-toric fibration, $\check{\varphi}$ and $\check{\varphi}_r$ agree up to compactly supported Hamiltonian isotopy. This readily follows from the two descriptions together with the discussion in Section 3.2.2 of the compatibility of the Hurwitz moves on the Lefschetz side and the nodal slides on the almost-toric side. \square

Remark 6.2. Recall the short exact sequence from Theorem 2.5:

$$1 \rightarrow N = \text{Hom}(\pi_1(Y_e \setminus D), \mathbb{C}^*) \rightarrow \text{Aut}(Y_e, D; \text{cpt}) \rightarrow \text{Adm}/W \rightarrow 1.$$

We've just seen that for any element of Adm/W , we can construct a mirror (graded) symplectomorphism $\check{\varphi}$. The autoequivalence of the Fukaya category mirror to an element of N is clear: it is given by the change of local systems encoded by the corresponding element of $\text{Hom}(\pi_1(M), \mathbb{C}^*)$.

Remark 6.3. On the Lefschetz side, we could construct the M_t and $\check{\varphi}_t$, $t \in [0, r]$, by deforming the base of the Lefschetz fibration by dragging around critical values in the base in the way prescribed by the Hurwitz moves. See Figure 6.1 for a single nodal slide.




FIGURE 6.1. Deforming the base of the Lefschetz fibration to get the isotopy from M to M_1 .

6.2. Independence of choices. We want to prove that the map $\check{\varphi}$ constructed in Theorem 6.1 is independent of choices. In order to do this, we start by understanding the mirrors to $SL_2(\mathbb{Z}) \subset \text{Aut}(\mathbb{C}^*)^2 \subset \text{Bir}((\mathbb{C}^*)^2)$.

6.2.1. Mirrors to $SL_2(\mathbb{Z})$ transformations. The $SL_2(\mathbb{Z})$ action on $(\mathbb{C}^*)^2$ can be described as follows. The B side is standard: for a given arbitrary toric pair (\bar{Y}, \bar{D}) , any element g of $SL_2(\mathbb{Z})$ gives a birational transformation of \bar{Y} . After sufficiently many toric blow ups, say to (\bar{Y}', \bar{D}') , g lifts to a biholomorphism from \bar{Y}' to, say, \bar{Y}'' . If (\tilde{Y}', \tilde{D}') is given by interior blow-ups on (\bar{Y}', \bar{D}') in m_1, \dots, m_k points, then g induces an automorphism from (\tilde{Y}', \tilde{D}') to $(\tilde{Y}'', \tilde{D}'')$, given by blowing up $(\tilde{Y}'', \tilde{D}'')$ according to the cyclic permutation of the m_i determined by g . (If g preserves the fan for (\bar{Y}', \bar{D}') and the m_i are cyclically symmetric, we get an automorphism of (\tilde{Y}', \tilde{D}') to itself.)

On the A side, let M be the mirror symplectic manifold to (\tilde{Y}', \tilde{D}') , and let M' be the mirror to $(\tilde{Y}'', \tilde{D}'')$. By assumption, we can set up the almost-toric fibrations on M and M' so that one base is mapped to the other by g , with critical points and invariant lines mapped to each other. This naturally lifts to a symplectomorphism from M to M' , say \check{g} .

By using suitable Liouville forms, one could arrange for \check{g} to induce a map from $\mathcal{W}(M)$ to $\mathcal{W}(M')$. If we knew that there was a compatible map of directed Fukaya categories, Proposition

2.14 would apply to show that \check{g} corresponds to g under homological mirror symmetry. (Note that this would not require a fine-level description of the action on those categories!) This should follow from [HW], who consider a version of the directed Fukaya category defined in terms of sections of the almost-toric fibration. In the current paper, we only use this for conjugates of compactly supported maps by \check{g} , which as usual follows from Proposition 2.14 together with Propositions 5.1 and 5.2.

Lemma 6.4. *Let g and \check{g} be as above. Assume that $\check{h} \in \pi_0 \text{Sym}_c^{\text{gr}} M$ is a symplectomorphism of M mirror to an autoequivalence $h \in \text{Auteq } D(Y)$. Then $\check{g}^{-1} \circ \check{h} \circ \check{g} \in \pi_0 \text{Sym}_c M$ is mirror to $g^{-1} \circ h \circ g \in \text{Auteq } D(Y)$.*

Remark 6.5. Alternatively, to get a well-defined action on directed categories, we could set up a mirror \tilde{g} to g in terms of the Lefschetz fibration by combining the almost-toric / Lefschetz identification with [HK, Proposition 3.19]. The map \tilde{g} would be a fibred symplectomorphism constructed by combining a rotation of the fibre near infinity with an automorphism of the base of w (similarly to the proof of Theorem 6.1). While this is less geometrically appealing, such a map \tilde{w} will by construction automatically act on $D^b \mathcal{F}^\rightarrow(w)$, and be mirror to $g \in \text{Auteq } D(Y)$ under our HMS correspondence. While conditions nears infinity are different for \tilde{g} and \check{g} (heuristically, they are suited for different versions of the directed Fukaya category), they will agree on arbitrarily large compact sets. (In particular, they will be interchangeable for the purposes of conjugating a compactly supported map.)

6.2.2. Proof of independence of choices.

Proposition 6.6. *The symplectomorphism $\check{\varphi}$ constructed in Theorem 6.1 is independent of choices.*

In order to prove this, it is enough to show that two different factorisations of φ into elementary transformations induce mirror symplectomorphisms which agree up to a compactly supported Hamiltonian isotopy. (This also takes care of the choice of toric model.) This means that we need to check that relations in the group of birational automorphisms of $(\mathbb{C}^*)^2$ induce isotopic symplectomorphisms. We start with the most interesting one.

The ‘ A_2 cluster relation’. There is a well-known order five relation in the group of birational transformations of $(\mathbb{C}^*)^2$, which experts will recognise as corresponding to the fact that there are exactly five cluster charts for the A_2 quiver. To realise this relation (minimally) in terms of automorphisms of a log CY pair, take the model $(\tilde{Y}, \tilde{D}) \rightarrow (\bar{Y}, \bar{D})$ where (\bar{Y}, \bar{D}) is a corner blow-up of $\mathbb{P}^1 \times \mathbb{P}^1$, and (\tilde{Y}, \tilde{D}) is given by one interior blow up on each of the self-intersection zero components. (This is a compactification of the A_2 cluster variety.) The mirror almost-toric fibration is given in the left-hand side of Figure 6.2; label the boundary components as in the




FIGURE 6.2. Almost toric fibration for the ‘ A_2 quiver relation’: initial configuration, and mirrors to E and then g . The cuts are in black, and dotted lines correspond to fan vectors (including ones with no interior blow-ups).

figure. (Both the A and B side spaces, and the sequences of mutations, are carefully described in [CV].)

Let E be the elementary transformation at \bar{D}_5 , and let g be the $SL_2(\mathbb{Z})$ birational transformation which takes \bar{D}_i to \bar{D}_{i-1} . Let $P = gE$; this defines an automorphism of \tilde{Y} (rather than a mere birational transformation), permuting the components of D ; we have $P^5 = \text{Id}$. Equivalently, setting $E_a = g^{-a}Eg^a$, we have $E_4 \dots E_0 = \text{Id}$ (note $g^{-4} = g$). On the symplectic side, we have the following.

Proposition 6.7. *Let $(\tilde{Y}, \tilde{D}) \rightarrow (\bar{Y}, \bar{D})$ be as above, and let M be its mirror. Let $\tilde{\varphi}$ be the compactly supported symplectomorphism of M associated to the factorisation of $\text{Id} \in \text{Aut}(\tilde{Y})$ into $E_4 \dots E_0$; then $\tilde{\varphi}$ is isotopic to the identity through a compactly supported Hamiltonian.*

Symplectic incarnations of the five A_2 cluster charts have been extensively studied, going back to [EHK16] – see e.g. [Pan17, STWZ19, TZ18, GSW, STW, CV]. Nevertheless, Proposition 6.7 doesn’t readily follow from the existing literature, as the focus has instead largely been on proving that symplectic objects ‘mirror’ to the five charts, such as Lagrangian fillings of the right-hand trefoil, are distinct. (Note that by Theorem 6.1 we already know that $\tilde{\varphi}$ acts as the identity on the Fukaya category.) We instead give a self-contained proof of Proposition 6.7, and comment further on the relation with the contact geometry literature in Remark 6.8.

Proof. of Proposition 6.7. The isotopy between $\tilde{\varphi}$ and the identity is hard to see directly from the perspective of the almost-toric fibration. We work instead with the Lefschetz fibration $w : M \rightarrow \mathbb{C}$.

As described in Theorem 6.1, $\tilde{\varphi}$ is a fibred symplectomorphism with respect to w ; as before, let $f_{\tilde{\varphi}}$ be the automorphism of the base. We calculate this by hand. The key is to use, first, a more symmetric description of w , given in Figure 6.3.

We can now iteratively apply the isotopy corresponding to a single elementary transformation (in Figure 6.1), combined with [HK, Proposition 3.19] for cycling the labels on components of \bar{D} , to get a description of $f_{\tilde{\varphi}}$; this is carried out in Figure 6.4.

The upshot is given in Figure 6.5: $f_{\tilde{\varphi}}$ is given by five half-twists between the critical points for W_5 and $\tau_{V_0}^{-1}W_1$. On the other hand, one can immediately check that these two vanishing




FIGURE 6.3. Lefschetz fibration $w : M \rightarrow \mathbb{C}$: standard (left), cyclically symmetric viewpoint (center), and in relation to the first mutation (same colour-coding as Figure 3.2).




FIGURE 6.4. Calculating f_φ by applying the isotopies corresponding to ξ_0, \dots, ξ_4 . We’re using the same cyclically symmetric configuration as in Figure 6.3: T is given by gluing the blue and green thimbles along the purple arc.

cycles are in a local A_2 configuration; in particular, we can deform w so that these two critical points merge (along their vanishing paths) to give a single A_2 critical point, at which point the half-twists readily unravel to give an isotopy to the identity. (Alternatively, as the sub-Lefschetz fibration on which φ has support is just a copy of the ‘standard’ A_2 fibration, with total space \mathbb{C}^2 , one could quote [Gro85].) \square

Remark 6.8. Intuitively, the key to Proposition 6.7 is to show that $\varphi(T)$ is Hamiltonian isotopic to T . This can be checked using tools from contact geometry, which we now sketch. In fact, a slightly stronger statement is true: we’ll see that the statement holds relative to




FIGURE 6.5. Upshot: $f_{\tilde{\varphi}}$ is given by five half-twists between two vanishing cycles

a disc on T (fixed pointwise under the five mutations and then the Hamiltonian isotopy), or, equivalently, we can work with a once-punctured torus. First, recall that the max-tb right-handed Legendrian trefoil knot $K \subset S^3 = \partial(B^4, \omega_0)$ has at least five exact Lagrangian fillings, all once-punctured tori ([EHK16, Example 8.4], see also [Pan17, TZ18]). On the other hand, the Weinstein domain given by attaching a Weinstein 2-handle along K is precisely M – see e.g. [CM19, Section 4.1], or the later conceptual argument in [Cas, Corollary 1.2]. One can then proceed, for instance, using the framework of weaves introduced in [CZ]: first, the five Lagrangian fillings can be described in terms of 2-weaves, in this case trivalent graphs, see [CZ, Section 7.1.3]; the fillings are given by the 2-weaves dual to the five triangulations of the pentagon. Second, Lagrangian mutations can be understood in that language: they are given by Whitehead moves on the trivalent graph, dual to flips in the triangulation, see [CZ, Theorem 4.21]. Applying this, we get that starting with any of our five Lagrangian fillings and performing the sequence of five mutations gives a filling which is Hamiltonian isotopic, relative to the boundary K , to the filling we started with.

We can now conclude:

Proof. of Proposition 6.6. Relations for the group of birational transformations of $(\mathbb{C}^*)^2$ were described by Blanc in [Bla13, Theorem 2]. In our case, they consist of, first, relations in $SL_2(\mathbb{Z})$. These are readily satisfied on the symplectic side: the conjugate by $g \in SL_2(\mathbb{Z})$ of an elementary transformation with respect to ray v_j is the elementary transformation with respect to ray $g \cdot v_j$, and the mirror statement is automatic. Second, Blanc’s list gives us two relations involving elementary transformations which need checking on the symplectic side. The first one ($PCP = I$ in his notation) amounts to checking that mutating and then mutating back gives the identity, which is automatic geometrically; and the second relation ($P^5 = 1$ in his notation) was covered by Proposition 6.7. \square

6.3. Examples. We calculate the first few explicit examples of nodal slide recombinations.

Example 6.9. *B_2 cluster variety relation.* Consider the mirror M to the pair (Y, D) given in Figure 6.6. (It has $k = 6$, with three interior blow-ups; experts will recognise a standard

compactification of the B_2 cluster variety, see [CV].) M has a non-compactly supported symplectomorphism ρ described in the figure; this consists of a double nodal slide and cut transfer, followed by a single one, followed by an $SL_2(\mathbb{Z})$ map. By construction, ρ^3 has compact support; as $\text{Adm}/W = \{1\}$ in this case, ρ^3 is Hamiltonian isotopic to the identity by Theorem 6.1. This gives a symplectic mirror to what is sometimes known as the ‘ B_2 cluster variety relation’.




FIGURE 6.6. Nodal slides for Example 6.9; the SL_2 action is given by taking the i th ray on the right-hand side to the $i - 2$ th, with indices mod 6.

Assume that (Y, D) is negative semi-definite, so that Y is a rational elliptic surface, and $M = X \setminus E$ the complement of a smooth anticanonical divisor in a del Pezzo surface. From Section 2.2.2, we know that $\text{Auteq}(Y, D; \text{pt})$ is infinite in two cases, and, in both, equal to $k \cdot MW(Y, \mathbb{P}^1)$, where $MW(Y, \mathbb{P}^1) \simeq \mathbb{Z}$ is the Mordell-Weil group of Y . For both of these cases, we can factorise a generator for $MW(Y)$ and write down the mirror sequence of nodal slides, together with the $SL_2(\mathbb{Z})$ action corresponding to the permutation of the components of D .

Example 6.10. *Mirror to Mordell-Weil for $\text{Bl}(\mathbb{P}^1 \times \mathbb{P}^1) \setminus E$.* Assume that $k = 7$; X is the blow-up of $\mathbb{P}^1 \times \mathbb{P}^1$ in a point. A generator φ for $MW(Y, \mathbb{P}^1)$ gives the symplectomorphism $\check{\varphi}$ of M described by the sequence of nodal slides in Figure 6.8, postcomposed by the $SL_2(\mathbb{Z})$ action encoded by the labeling of the invariant rays. The map $\check{\varphi}^7$ has compact support, and, by Theorem 6.1, is mirror to φ^7 .




FIGURE 6.7. Nodal slides for Example 6.10; the SL_2 action is determined by taking to i th ray in the right-hand side diagram to the $i + 1$ th.

Example 6.11. *Mirror to Mordell-Weil for $\mathbb{F}_1 \setminus E$.* Assume that $k = 8$ and $X = \mathbb{F}_1$. A generator φ for $MW(Y, \mathbb{P}^1)$ gives the symplectomorphism $\check{\varphi}$ of M described by the sequence of nodal slides in Figure 6.8, postcomposed by the $SL_2(\mathbb{Z})$ action encoded by the labeling of the invariant rays. The map $\check{\varphi}^8$ has compact support, and, by Theorem 6.1, is mirror to φ^8 .




FIGURE 6.8. Nodal slides for Example 6.11; the SL_2 action is determined by taking to i th ray in the right-hand side diagram to the $i + 1$ th.

6.4. Relations between different families of symplectomorphisms. Homological mirror symmetry immediately hands us symplectic versions of Lemma 2.15 at the level of Fukaya categories. The first two relations automatically have geometric counterparts: given a Lagrangian sphere S and a symplectomorphism f , $\tau_{f(S)} = f \circ \tau_S \circ f^{-1} \in \pi_0 \text{Sym}_c^{\text{gr}} M$. We check the other two.

Lemma 6.12. *Suppose that $\sigma_L \in \pi_0 \text{Sym}_c^{\text{gr}} M$ is the Lagrangian translation associated to $[L - L_0] \in H_2(M)/H_2(\partial M)$ and $\tilde{\varphi} \in \pi_0 \text{Sym}_c^{\text{gr}} M$ a nodal slide recombination. Then*

$$\sigma_{\varphi_* L} = \varphi \circ \sigma_L \circ \varphi^{-1} \in \pi_0 \text{Sym}_c M.$$

Similarly with non-compactly supported Lagrangian translations.

Proof. This follows from iteratively applying Lemma 4.13. Similarly with non-compactly supported Lagrangian translations. \square

Suppose $C \subset Y \setminus D$ is a (-2) curve and $a \in \mathbb{Z}$. Recall that as autoequivalences of derived categories,

$$T_{\mathcal{O}_C(a-1)} \circ T_{\mathcal{O}_C(a)} = _ \otimes \mathcal{O}_Y(C).$$

We will prove a symplectic analogue of this relation in Lemma 6.13. Let's start with the set-up.

Choose an almost-toric fibration $\pi : M \rightarrow \mathbb{R}^2$ as in Proposition 5.2, and let $s_a = \sigma_E^{a+1} s_C$ be the sphere mirror to $i_* \mathcal{O}_C(a)$ constructed therein; these all project to a segment γ between two critical points of π . This fibration comes with a preferred reference Lagrangian section L_0 ; using this, let σ_{L_C} be the Lagrangian translation mirror to tensoring with $\mathcal{O}_Y(C)$ from Theorem 5.5.

Let $B_{\text{loc}} \subset \mathbb{R}^2$ be a small open neighbourhood of γ , and let $M_{\text{loc}} = \pi^{-1}(B_{\text{loc}})$. (This should be thought of as mirror to the formal neighbourhood of a (-2) curve, or to the resolution of an A_1 singularity, cf. [CU13].) M_{loc} is the total space of a Lefschetz fibration well-known to experts, which we briefly recall. First, consider

$$M'_{\text{loc}} = \{xy + (z-1)(z-2) = 0\} \subset \mathbb{C}^2 \times \mathbb{C}^*$$

equipped with the Kaehler form with potential $|x|^2 + |y|^2 + (\log |z|)^2$. Let $f : M'_{\text{loc}} \rightarrow \mathbb{C}^*$ be projection to z , a conic Lefschetz fibration with two critical fibres. There is a symplectic S^1 action given by $(x, y, z) \mapsto (e^{i\theta}x, e^{-i\theta}y, z)$. Following [Aur07, Section 5.1], let $\delta_{z_0}(x_0, y_0)$ be the signed symplectic area between the equator $\{|x| = |y|\} \subset f^{-1}(z_0)$ and the S^1 orbit of (x_0, y_0, z_0) . Then the map $\pi' : M'_{\text{loc}} \rightarrow \mathbb{R}^2$ given by $\pi'(x, y, z) = (|z|, \delta_z(x, y))$ defines a singular Lagrangian torus fibration: symplectic parallel transport with respect to f is S^1 -equivariant and preserves δ . There are two nodal critical fibres, the preimages of $(1, 0)$ and $(2, 0)$, and one then readily gets that after suitably truncating, M'_{loc} agrees with M_{loc} . Moreover, the spheres s_a can be represented as matching cycles with respect to f , as in Figure 6.9.




FIGURE 6.9. Lagrangian spheres s_a : matching paths in the base of the Lefschetz fibration $f : M'_{\text{loc}} \rightarrow \mathbb{R}^2$. The matching path for s_a can be obtained from the one for s_{a-1} by applying a full negative twist in the segment from 0 to 1.

Lemma 6.13. *Suppose that we are given $\sigma_{LC} \in \pi_0 \text{Symp}_c(M)$ and Lagrangian spheres s_a , $a \in \mathbb{Z}$ as above. Then*

$$\tau_{s_{a-1}} \circ \tau_{s_a} = \sigma_{LC} \in \pi_0 \text{Symp}_c(M).$$

Proof. It's enough to prove this for $a = 0$. Set $\rho = \tau_{s_{-1}} \circ \tau_{s_0}$. The local model for the Dehn twist in the matching cycle s_a is well known: it is induced by the half-twist tw_a in the associated matching path (as an isotopy class of automorphism of \mathbb{C}^* relative to the marked points), see e.g. [Sei01, Sei03, AMnP05]. In the case at hand, recall that we have a two-parameter family of Lagrangian tori, with each one fibred over one of the concentric circles $\eta_c = \{|z| = c\}$, for some $c \in \mathbb{R}_+$. The family $\{\eta_c\}$ foliates the base of the Lefschetz fibration f . On the other hand, the image of the foliation under $\text{tw}_{-1} \circ \text{tw}_0$ is isotopic relative to marked points to the original one. See Figure 6.10. (This is familiar to experts, see e.g. [Sei, Remark 11.11] for the observation for a single η_c .) This implies that after post-composing with a Hamiltonian isotopy, we can arrange for $\tau_{s_{-1}} \circ \tau_{s_0}$ to respect the almost-toric fibration $M_{\text{loc}} \rightarrow B_{\text{loc}}$. The Arnol'd–Liouville theorem then strongly constrains the map: it must be the Lagrangian translation determined by the image of a reference section L_0 (the argument extends over the nodal fibres similarly to the proof of Proposition 4.9). Reading off the action on homology is then enough to identify this translation with σ_{LC} . \square

7. APPLICATIONS

7.1. New symplectomorphisms. Fix a log CY surface (Y_e, D) with mirror M . Let \bar{Q} , Adm and W be as before, and let \bar{Q}_{symp} , $\text{Adm}/W_{\text{symp}}$ and Br_{symp} be the subgroups of




FIGURE 6.10. Action of $tw_{-1} \circ tw_0$ on the base of the Lefschetz fibration $f : M'_{loc} \rightarrow \mathbb{R}^2$, with the images of s_0 , s_{-1} , and a sample of circles γ_c .

$\pi_0 \text{Symp}_c^{\text{gr}}(M)$ generated by respectively, compactly supported Lagrangian translations, nodal slide recombinations, and Dehn twists in arbitrary Lagrangian spheres. (For the latter we do not restrict ourselves to the ‘known’ spheres constructed in Proposition 5.2.)

Theorem 7.1. *Given the above setting, we have the following:*

- (1) *The map $\bar{Q} \times \text{Adm}/W \rightarrow \pi_0 \text{Symp}_c^{\text{gr}}(M)$ with image $\bar{Q}_{\text{symp}} \times \text{Adm}/W_{\text{symp}}$ is faithful.*
- (2) *Let $\langle \Phi \rangle \leq \bar{Q}$ be the sublattice generated by classes of roots. Then*

$$\text{Br}_{\text{symp}} \cap (\bar{Q}_{\text{symp}} \times \text{Adm}/W_{\text{symp}}) = \langle \Phi \rangle_{\text{symp}} \times \{1\}$$

In particular, whenever $\text{Adm} \neq W$ or $\bar{Q} \neq \langle \Phi \rangle$, we get elements of $\pi_0 \text{Symp}_c^{\text{gr}}(M)$ which are not products of Dehn twists.

Proof. (1). We have a well-defined geometric action by Lemma 6.12; faithfulness holds at the K -theoretic level.

(2). After the HMS identification, the K theory classes of Lagrangian spheres are the same as the ones in Remark 2.4. Also, recall that on the mirror side, $\text{Br}_Z \cap \text{Aut}(Y) = \{\text{Id}\} \in \text{Auteq}(Y)$. The claim then follows from mirrors of the relations between autoequivalences in Lemma 2.15, notably Lemma 6.13. \square

Remark 7.2. The classification in Remark 2.4 allows us to exclude arbitrary spherical twists, irrespective of whether they are given geometrically by Dehn twists in embedded Lagrangian spheres. Twists in other Lagrangians with periodic geodesic flow are ruled out for cohomological and dimension reasons. Less is known in general about the categorical action of fibred twists (see however [Per, WW16, MW18]). The only known fibred twists for our family of manifolds M are boundary twists in the semi-definite (simple elliptic) cases; these are given by a finite-time action of the Reeb flow on the boundary. In particular, they act trivially on relative homology, and so on $K(\mathcal{W}(M))$, and so do not lie in $\bar{Q}_{\text{symp}} \times \text{Adm}/W_{\text{symp}} \setminus \{\text{Id}\}$. (For hypersurface simple elliptic singularities, they are products of Dehn twists [Sei00, Section 4.c].)

Example 7.3. Nodal slide recombinations: Adm/W is non-trivial in the following cases.

- Small values of k : for $k \leq 6$, Adm/W is always trivial, and there are no interesting nodal slide recombinations. On the other hand, for $k = 7$, Adm/W is always infinite; this is also true when $k = 8$ and $\pi_1(U) = 0$.
- In general, note that if Adm_Y/W_Y is non trivial for a log CY pair (Y, D) , and a log CY pair (Y', D') is obtained from (Y, D) through any sequence of toric and interior blow-ups, then $\text{Adm}_{Y'}/W_{Y'}$ is also non trivial – indeed, pullback gives an injection $\text{Adm}_Y/W_Y \rightarrow \text{Adm}_{Y'}/W_{Y'}$. In particular, the examples for small values of k can be used to generate many more.
- When $Y = Y_e$, $\text{Aut}(Y_e)$ acts with rational polyhedral domain on the nef cone [Li]. This implies that Adm/W is infinite if and only if the nef cone is not rational polyhedral.

Example 7.4. \bar{Q} is not typically generated by classes of simple roots. Some specific examples include:

- Semi-definite cases with D a cycle of (-2) curves and $k \geq 7$, except when $k = 8$ and $\pi_1(M) = \mathbb{Z}/2$.
- Whenever the rank of Q is one and D is negative (semi-)definite.
- Examples 4.3 and 4.4. in [Fri13]. Both have the property that Q has rank 2 and the root system is empty.

Remark 7.5. In the semi-definite case, a compactly supported symplectomorphism of M induces a symplectomorphism of its compactification X , a del Pezzo surface with Kaehler anti-canonical form. However, in our cases the induced symplectomorphisms of X act trivially on homology, and thus are isotopic to the identity [Gro85, LP04, Eva11].

7.2. Lagrangian spheres and symplectic Torelli group. For small values of k , a lot is explicitly known about (-2) curves in $Y \setminus D$; as soon as a toric model for (Y, D) involves two interior blow ups on the same component of \bar{D} , there is at least one (-2) curve. Moreover, suppose that D is negative definite and does not contain a (-1) curve. Then for $k \leq 5$, there are finitely many (-2) curves on Y_e , which give a basis of Q [Loo81]. For $k = 6$, the (-2) curves span Q , with exactly one relation between them; in particular, there's also finitely many of them [Li, Sim]. For $k = 7$, there are always infinitely many (-2) curves; for $k = 8$ there are infinitely many whenever the fundamental group of the complement U_e is trivial. (For $k \geq 9$ little has been done systematically, though individual cases can be done by hand.)

7.2.1. Infinitely many (-2) curves. Assume that $Y \setminus D$ contains infinitely many (-2) curves (necessarily countable). This implies we are in the negative definite case.

Corollary 7.6. *Let $(Y, D) = (Y_e, D)$ be a log Calabi-Yau surface with distinguished complex structure, and let $w : M \rightarrow \mathbb{C}$ be its mirror, i.e. a Milnor fibre of the dual cusp of D . Assume that Y contains infinitely many (-2) curves. Then:*

- (1) *There is a countable infinite collection of Lagrangian spheres in M , none of which is contained in the subcategory of $D^b\mathcal{W}(M)$ split-generated by the others.*
- (2) *The Dehn twist in any one of them is not contained in the subgroup of $\pi_0 \text{Symp}_c(M)$ generated by any of the others; similarly with squares of Dehn twists.*

Proof. Let C_i , $i \in \mathbb{N}$, be an infinite collection of (-2) curves in Y ; wlog they all lie in U . Let $S_i = s_{C_i}$ be the Lagrangian sphere mirror to $i_*\mathcal{O}_{C_i}(-1)$, from Proposition 5.2. Let $p_i \in C_i \setminus \cup_{j \neq i} C_j$. By considering morphisms with $i_*\mathcal{O}_{p_i}$, we see that $i_*\mathcal{O}_{C_i}(-1)$ cannot be contained by the subcategory split-generated by $\{i_*\mathcal{O}_{C_j}(-1)\}_{j \neq i}$. The homological mirror symmetry theorem then implies the first point. For the second one, one can proceed similarly, noting that $i_*\mathcal{O}_{p_i}$ is fixed by the spherical twist in $i_*\mathcal{O}_{C_j}(-1)$ for $j \neq i$, but not under the twist in $i_*\mathcal{O}_{C_i}(-1)$, nor its square. \square

Remark 7.7. Using [SS20a], the same statements and proofs hold for K3 surfaces whose mirror K3s containing infinitely many embedded (-2) curves.

Discussion. All of the spheres in Corollary 7.6 can be thought of as vanishing cycles for the cusp singularity dual to the one with cycle D , with Dehn twists in them induced by monodromy maps in the moduli spaces from Section 1.5. In particular, Corollary 7.6 should be contrasted with the case of (arbitrary) hypersurface singularities, for which Picard–Lefschetz theory tells us that any distinguished collection of vanishing cycles generates all others as objects of the Fukaya category, and that Dehn twists in them generate the monodromy group. (Note that the full compact Fukaya category of M can never be split generated by spherical objects: this is immediate by using HMS and by considering supports of mirror objects.) On the other hand, by [Li], $\text{Aut}(Y)$ acts on the collection of (-2) curves with finitely many orbits (note that there’s no a priori uniform bound); thus there are finitely many orbits of known Lagrangian spheres in $D^b\mathcal{W}(M)$ under the action of compactly supported symplectomorphisms. There could be more than one orbit: for instance, in the $k = 6$ semi-definite case, Y_e is the total space of elliptic fibration where the boundary D is a cycle of 6 (-2) curves, and the interior (-2) curves form an I_2 and an I_3 fibre, giving two orbits of Lagrangian spheres in $D^b\mathcal{W}(M)$, where M is mirror to $Y_e \setminus D$. (Note that $\text{Adm} = W$ here.)

7.3. Semi-definite case: monodromy and mirror to the $SL_2(\mathbb{Z})$ action. Assume that D is negative semi-definite, i.e. that M is the Milnor fibre of a simple elliptic singularity. We re-visit the monodromy perspective from the introduction in this case. We have that $M = X_0 \setminus E_0$, where X_0 is a del Pezo surface, with E_0 smooth anticanonical. The natural

moduli space of complex structures for M consists of pairs (X, E) where X is a del Pezzo surface deformation equivalent to X_0 , and $E \subset X$ a smooth anticanonical divisor.

First, we want to carefully check that any loop in this moduli space induces a symplectomorphism of M , well-behaved near ∂M in the cases where it's not compactly supported. Assume we have a smooth loop of pairs $\{(X_t, E_t)\}_{t \in [0,1]}$ where the X_t are del Pezzo surfaces, and $E_t \subset X_t$ are smooth anticanonical elliptic curves. M is equipped with ω_0 , the restriction of a Kaehler form Poincaré dual to E_0 ; by making coherent choices of global sections of a smoothly varying very ample line bundle, we get a smooth loop of triples $\{(X_t, E_t, \omega_t)\}_{t \in [0,1]}$, where $\omega_t \in \Omega^2(X_t)$ is a Kaehler form such that $[\omega_t] = P.D[E_t] \in H^2(X_t; \mathbb{Z})$. Let $f_t : X_0 \rightarrow X_t$ parametrise the family. Set $M_t := X_t \setminus E_t$, and $M = M_0 = M_1$. By assumption, $\omega_t|_{M_t}$ is exact, and using the tubular neighbourhood theorem for symplectic submanifolds, we can choose a primitive θ_t for $\omega_t|_{M_t}$ such that on a collar neighbourhood of ∂M_t , the corresponding Liouville vector field radially scales each fibre of the symplectic normal bundle of E_t (inwards with respect to the deleted zero-section E_t). Moreover, after isotopy, we can assume that $f_t^{-1}(E_t) = E_0$, and that $f_t^* \theta_t = \theta_0$ in a neighbourhood of E_t ; note that f_1 can be arranged to restrict to the identity on this neighbourhood precisely when it induces the identity on $H_1(E_0; \mathbb{Z})$. A Moser argument gives a symplectomorphism $f : M \rightarrow M$, smoothly isotopic to f_1 rel boundary, which is the compactly supported whenever $(f_1)_*$ is the identity on $H_1(E_0; \mathbb{Z})$. Choosing a grading, the map f acts on the wrapped Fukaya category; using the framework of stops from [Syl19, GPS], we expect to arrange an action on the directed Fukaya category.

Consider $H_2(M; \mathbb{Z})$. The “tube over a cycle” map gives a primitive embedding $H_1(E; \mathbb{Z}) \subset H_2(M; \mathbb{Z})$. This sublattice can be characterised as the radical of the degenerate intersection form on $H_2(M; \mathbb{Z})$. Now suppose that g is an arbitrary symplectomorphism of M . It induces an automorphism of $H_2(M; \mathbb{Z})$ respecting the intersection form, and so restricts to an automorphism of $H_1(E; \mathbb{Z})$. If g is the result of a monodromy construction as above, we recover the original action on $H_1(E; \mathbb{Z})$. Dually, one could start with the action of g on $H_2(M, \partial M; \mathbb{Z})$, and get the (dual) action on $H^1(E; \mathbb{Z})$ via cap products. Alternatively, if g induces an autoequivalence of the directed Fukaya category of M , this could be obtained by looking at the induced element of $\text{Aut } K(D^b \mathcal{F}^\rightarrow(w))$ and then considering the action on the radical to get an element in $\text{Aut } H^1(E; \mathbb{Z})$. Call the resulting map $\tilde{\Theta} : \text{Auteq } D^b \mathcal{F}^\rightarrow(w) \rightarrow \text{Aut } H^1(E; \mathbb{Z})$. The map $\tilde{\Theta}$ is surjective: by considering paths of smooth elliptic curves in \mathbb{P}^2 (or $\mathbb{P}^1 \times \mathbb{P}^1$) and blowing up points on those curves, we can arrange to get arbitrary elements of $SL_2(\mathbb{Z})$ (cf. [Kea15, Section 5.4]).

Finally, recall that on the mirror side, we have the map $\Theta : \text{Auteq } D(Y) \rightarrow \text{End } K(F)$ of equation 2.13, where F denotes a smooth fibre of the elliptic fibration $\varpi : Y \setminus D \rightarrow \mathbb{C}$. $K(F)$ is generated by $[\mathcal{O}_{pt}]$ and $[\mathcal{O}_F]$. Now the radical of the intersection pairing on $H_2(Y \setminus D; \mathbb{Z})$ is the primitive sublattice $\langle \gamma, [F] \rangle$, where γ is the ‘linking torus’ for a node of D . Homological mirror symmetry induces an identification of this lattice with the sublattice $H_1(E; \mathbb{Z}) \subset H_2(M; \mathbb{Z})$; and by construction $\tilde{\Theta}$ is mirror to Θ . (On the B side, $K(F)$ should be thought of as dual to

$\langle \gamma, [F] \rangle = H_1(E; \mathbb{Z}) \subset H_2(M; \mathbb{Z})$, with γ corresponding to $[\mathcal{O}_{pt}]$ and F to $[\mathcal{O}_F]$. In particular, we expect that there is a non-compactly supported symplectomorphism mirror to the fibrewise Fourier–Mukai transform described in Section 2.3.2.

Remark 7.8. This should be compared with [Hic21, Section 5]: in the case where $X = \mathbb{CP}^2$, Hicks constructs a symplectomorphism g of $M = \mathbb{CP}^2 \setminus E$ (where E is smooth anticanonical), by using an automorphism of the Hesse pencil on \mathbb{CP}^2 , which interchanges an SYZ fibre and a tropical Lagrangian torus, L_{T^2} in his notation. The mirror log CY surface Y , denoted \check{X}_{9111} in [Hic21], is the total space of a rational elliptic fibration with an I_9 fibre above infinity and three other singular fibres, all nodal. [Hic21, Section 5] provides evidence that g is mirror to a fibrewise Fourier–Mukai transform on Y .

7.4. Semi-definite case: connection with the special Lagrangian viewpoint. We briefly discuss connections between our results and work of Collins–Jacob–Lin in the semi-definite case [CJLa, CJLb].

Suppose X is a del Pezzo surface, E a smooth anticanonical divisor, and $M = X \setminus E$. There exists a meromorphic volume form Ω on X with a simple pole along E ; this is unique up to scaling by a constant. Moreover, M has a complete Ricci flat metric [TY90], say g_{TY} . Let ω_{TY} be the associated Kaehler form (automatically exact).

Theorem 7.9. [CJLa] *Fix any primitive homology class $\alpha \in H_1(E, \mathbb{Z})$. Then M admits a special Lagrangian fibration $\pi_\alpha : M \rightarrow \mathbb{R}^2$ with respect to ω_{TY} , with special Lagrangian torus fibre the image of α under the tube map $H_1(E, \mathbb{Z}) \rightarrow H_2(M, \mathbb{Z})$.*

Note that once α has been specified, the (almost-)toric fibration underlying the special one should be unique up to nodal slides.

Let $(\alpha_\gamma, \alpha_F)$ be the basis of $H_1(E; \mathbb{Z})$ corresponding to $(\gamma, [F]) \subset H_2(Y \setminus U; \mathbb{Z})$ under the mirror identification. (As before, γ is the class of a linking torus for a node of D , and $[F]$ the class of a fibre of the elliptic fibration ϖ on the mirror Y_e .) Intuitively, exactly one of the fibrations above should be thought of as the SYZ fibration: the one π_γ with fibre in class $\text{Tube}(\alpha_\gamma) \in H_2(M; \mathbb{Z})$. Recall from Section 7.3 that monodromy considerations give non-compactly supported symplectomorphisms inducing any $SL_2(\mathbb{Z})$ action on $H_1(E; \mathbb{Z})$. We expect that for arbitrary α , π_α can be obtained as a Lagrangian fibration by precomposing π_γ with a suitable non-compactly supported symplectomorphism. In the case of π_{α_F} , this should be the mirror to the fibrewise Fourier–Mukai transform.

Collins–Jacob–Lin also study properties of the hyperkaehler structure associated to these Ricci flat metrics. Assume that the holomorphic volume form on M is normalised so that the special torus fibres have argument zero. Hyperkaehler rotation on M (from the I -structures to the J -structures with the standard conventions) yields a complex manifold U_{HK} , and π

induces a holomorphic fibration $\varpi_{\text{HK}} : U_{\text{HK}} \rightarrow \mathbb{C}$; by [CJLa, Theorem 6.4], U_{HK} admits a compactification to a relatively minimal elliptic fibration to \mathbb{P}^1 with an I_k fibre at infinity, where k is the degree of the del Pezzo surface X . In particular, this implies that Y is in the deformation component of the homological mirror Y_e to M .

While hyperkaehler rotation should generally *not* be thought of as mirror symmetry, an analysis in this case shows that the two can be set up to agree for a distinguished moduli space point: following [CJLb, Section 5] (and [McM07]), we can find a smooth del Pezzo surface X' deformation equivalent to X ; a smooth anticanonical elliptic curve $E \subset X'$; and a holomorphic volume form $\Omega^{M'}$ on M' such that:

- (1) Hyperkaehler rotation on M (from the I structures to the J ones) gives the distinguished complex structure $U_e = Y_e \setminus D$.
- (2) Under this hyperkaehler rotation, the special Lagrangian fibration on $M = X \setminus E$ with fibre class $\text{Tube}(\alpha_F)$, say π_F , becomes an elliptic fibration $\varpi^0 : U_e \rightarrow \mathbb{C}$ which compactifies to $\varpi : Y_e \rightarrow \mathbb{P}^1$.
- (3) The special Lagrangian fibration on M with fibre class $\text{Tube}(\alpha_\gamma)$ has fibres with argument $\pi/2$. After hyperkaehler rotation, this gives a special Lagrangian fibration on U_e .

Cf. [CJLb, Theorem 5.4]. This set-up can be used to obtain symplectomorphisms of M :

Lemma 7.10. [CJLb, Proposition 5.10] *Suppose we are in the setting above. Let $\psi \in \text{MW}(Y_e, \mathbb{P}^1)$ be an element of the Mordell-Weil group of Y . Then as a diffeomorphism of M , with the symplectic form given by hyperkaehler rotation, ψ is in fact a symplectomorphism of M , say ρ_ψ .*

Proof. This is essentially immediate from hyperkaehler identities. The map ψ is holomorphic on $U = U_e$ (whose complex structure is J), so it must at worst scale $\Omega_J = \Omega^U$. Moreover, the scaling factor must in fact be one: ψ acts by translation on each fibre, and uniformising the fibration near a smooth fibre one sees that the scaling factor must be one. (See [Huy16, Lemma 15.4.4] for details on an analogous argument, or the proof of [CJLb, Proposition 5.10] for a more analytic viewpoint.) Now split the equation $\psi^*\Omega_J = \Omega_J$ into real and imaginary parts to see that ψ preserves the symplectic form on M . \square

Suppose that ψ is a k th power in the Mordell-Weil group. The map ρ_ψ above doesn't have compact support: this is merely true asymptotically, in a sense which could be made precise. However, we observe that the ideas of Section 4 can be used in this setting to improve this. We first note the following:

Lemma 7.11. *Let s_0 be the choice of reference holomorphic section for the group law in $\text{MW}(Y_e, \mathbb{P}^1)$, and s_ψ the one for ψ . Then their images under hyperkaehler rotation are*

Lagrangian sections of π_F , say L_e and L_ψ , respectively. Moreover, away from critical fibres (and on fibres with a single critical point), ρ_ψ is precisely equal to the Lagrangian translation of $\pi_\varpi : M \rightarrow \mathbb{R}^2$ which takes L_e to L_ψ .

Proof. Hyperkaehler identities imply that holomorphic sections of ϖ become Lagrangian sections of π_F . By Lemma 7.10, the map ρ_ψ is a symplectomorphism of M , preserving the fibres of π_ϖ . By [Sym03, Lemma 2.5], this implies that on any Arnol'd–Liouville chart $V \times T^2$, $V \subset \mathbb{R}^2$, it is of the form $(q, p) \mapsto (q, p + F(q))$, where F is the derivative of a smooth function $V \rightarrow \mathbb{R}$. The claim is then immediate. \square

Corollary 7.12. *Let $\psi \in k \cdot MW(Y_e, \mathbb{P}^1)$ be an element of the Mordell–Weil group of Y_e fixing D pointwise. Then ρ_ψ is Hamiltonian isotopic, via a fibre-preserving isotopy of $\pi_F : M \rightarrow \mathbb{R}^2$, to a compactly supported symplectomorphism, say ρ_ψ^c .*

Proof. As $\psi \in k \cdot MW(Y_e, \mathbb{P}^1)$, the two sections s_0 and s_ψ defining ψ intersect at infinity, at an interior point of a component of $D = \varpi^{-1}(\infty)$. Thus the sections s_0 and s_ψ are smoothly isotopic near infinity; hence L_e and L_ψ are too. It then follows from Corollary 4.8 after a fibre-preserving Hamiltonian isotopy L_e and L_ψ can be taken to agree outside a compact set. (Recall the obstruction is the symplectic area of an annulus; one observes that it can be capped off with two Lagrangian discs, given by L_e and L_ψ , to get a closed chain; and as $\omega \in \Omega^2(M)$ is exact, the obstruction must thus vanish.) Lagrangian translation between L_e and the images of L_ψ under isotopy then gives the desired isotopy of ρ_ψ to a compactly supported symplectomorphism. \square

As before, this gives non-trivial symplectomorphisms for two cases: $k = 7$, or $k = 8$ with $X = \mathbb{F}_1$. By construction, the K -theory actions of ψ and ρ_ψ^c are mirror. For the $k = 8$ case, as there are no interior (-2) curves in Y_e , Proposition 2.14 immediately implies that the maps are HMS mirrors. (For the $k = 7$ case, there is an interior I_2 fibre, and no a priori reason for the two maps not to differ by a Torelli autoequivalence generated by twists in the associated spherical objects.)

REFERENCES

- [ABC⁺09] Paul S. Aspinwall, Tom Bridgeland, Alastair Craw, Michael R. Douglas, Mark Gross, Anton Kapustin, Gregory W. Moore, Graeme Segal, Balázs Szendrői, and P. M. H. Wilson. *Dirichlet branes and mirror symmetry*, volume 4 of *Clay Mathematics Monographs*. American Mathematical Society, Providence, RI; Clay Mathematics Institute, Cambridge, MA, 2009. [4.1](#), [4.4](#), [4.1](#)
- [AKO06] Denis Auroux, Ludmil Katzarkov, and Dmitri Orlov. Mirror symmetry for del Pezzo surfaces: vanishing cycles and coherent sheaves. *Invent. Math.*, 166(3):537–582, 2006. [2.8](#), [3.2](#)
- [AMnP05] Denis Auroux, Vicente Muñoz, and Francisco Presas. Lagrangian submanifolds and Lefschetz pencils. *J. Symplectic Geom.*, 3(2):171–219, 2005. [6.4](#)

- [Arn89] V. I. Arnol'd. *Mathematical methods of classical mechanics*, volume 60 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, [1989?]. Translated from the 1974 Russian original by K. Vogtmann and A. Weinstein, Corrected reprint of the second (1989) edition. [4.1](#)
- [Arn95] V. I. Arnol'd. Some remarks on symplectic monodromy of Milnor fibrations. In *The Floer memorial volume*, volume 133 of *Progr. Math.*, pages 99–103. Birkhäuser, Basel, 1995. [1.1](#)
- [Aur07] Denis Auroux. Mirror symmetry and T -duality in the complement of an anticanonical divisor. *J. Gökova Geom. Topol. GGT*, 1:51–91, 2007. [1.2](#), [3.2.2](#), [6.4](#)
- [BHPVdV04] Wolf P. Barth, Klaus Hulek, Chris A. M. Peters, and Antonius Van de Ven. *Compact complex surfaces*, volume 4 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, second edition, 2004. [2.2.2](#)
- [Bla13] Jérémy Blanc. Symplectic birational transformations of the plane. *Osaka J. Math.*, 50(2):573–590, 2013. [1.2](#), [3.2.1](#), [6.2.2](#)
- [Bri98] Tom Bridgeland. Fourier-Mukai transforms for elliptic surfaces. *J. Reine Angew. Math.*, 498:115–133, 1998. [2.3.2](#)
- [BS10] Tom Bridgeland and David Stern. Helices on del Pezzo surfaces and tilting Calabi-Yau algebras. *Adv. Math.*, 224(4):1672–1716, 2010. [5.1](#)
- [Cas] Roger Casals. Lagrangian Skeleta and Plane Curve Singularities. arXiv:2009.06737. [6.8](#)
- [CDRGG] Baptiste Chantraine, Georgios Dimitroglou Rizell, Paolo Ghiggini, and Roman Golovko. Geometric generation of the wrapped Fukaya category of Weinstein manifolds and sectors. arXiv:1712.09126. [4.3](#)
- [CFP] Alessio Corti, Matej Filip, and Andrea Petracci. Mirror Symmetry and smoothing Gorenstein toric affine 3-folds. arXiv:2006.16885. [5.1](#)
- [CJLa] Tristan Collins, Adam Jacob, and Yu-Shen Lin. Special lagrangian submanifolds of log Calabi-Yau manifolds. arXiv:1904.08363. [7.4](#), [7.9](#), [7.4](#)
- [CJLb] Tristan Collins, Adam Jacob, and Yu-Shen Lin. The SYZ mirror symmetry conjecture for del Pezzo surfaces and rational elliptic surfaces. arXiv:2012.05416. [7.4](#), [7.4](#), [7.4](#), [7.10](#), [7.4](#)
- [CK16] Alessio Corti and Anne-Sophie Kaloghiros. The Sarkisov program for Mori fibred Calabi-Yau pairs. *Algebr. Geom.*, 3(3):370–384, 2016. [1.2](#), [1.5](#)
- [CL09] Serge Cantat and Frank Loray. Dynamics on character varieties and Malgrange irreducibility of Painlevé VI equation. *Ann. Inst. Fourier (Grenoble)*, 59(7):2927–2978, 2009. [2.2.1](#)
- [CM19] Roger Casals and Emmy Murphy. Legendrian fronts for affine varieties. *Duke Math. J.*, 168(2):225–323, 2019. [6.8](#)
- [CU13] Kwokwai Chan and Kazushi Ueda. Dual torus fibrations and homological mirror symmetry for A_n -singularities. *Commun. Number Theory Phys.*, 7(2):361–396, 2013. [6.4](#)
- [CV] Man-Wai Mandy Cheung and Renato Vianna. Algebraic and symplectic viewpoint on compactifications of two-dimensional cluster varieties of finite type. arXiv:2008.03265. [6.2.2](#), [6.2.2](#), [6.9](#)
- [CZ] Roger Casals and Eric Zaslow. Legendrian Weaves: N-graph Calculus, Flag Moduli and Applications. arXiv:2007.04943. [6.8](#)
- [Dui80] J. J. Duistermaat. On global action-angle coordinates. *Comm. Pure Appl. Math.*, 33(6):687–706, 1980. [4.1](#)
- [EF21] Philip Engel and Robert Friedman. Smoothings and rational double point adjacencies for cusp singularities. *J. Differential Geom.*, 118(1):23–100, 2021. [1.2](#), [1.11](#)
- [EHK16] Tobias Ekholm, Ko Honda, and Tamás Kálmán. Legendrian knots and exact Lagrangian cobordisms. *J. Eur. Math. Soc. (JEMS)*, 18(11):2627–2689, 2016. [6.2.2](#), [6.8](#)

- [Eng18] Philip Engel. Looijenga’s conjecture via integral-affine geometry. *J. Differential Geom.*, 109(3):467–495, 2018. [1.1](#)
- [Eva11] Jonathan David Evans. Symplectic mapping class groups of some Stein and rational surfaces. *J. Symplectic Geom.*, 9(1):45–82, 2011. [1.1](#), [7.5](#)
- [FG09] V. V. Fock and A. B. Goncharov. Cluster ensembles, quantization and the dilogarithm. II. The intertwiner. In *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. I*, volume 269 of *Progr. Math.*, pages 655–673. Birkhäuser Boston, Boston, MA, 2009. [1.3](#)
- [Fri] Robert Friedman. On the geometry of anticanonical pairs. arXiv:1502.02560. [2.1](#), [2.2](#), [2.2.1](#), [2.7](#), [2.2.2](#)
- [Fri13] Robert Friedman. On the ample cone of a rational surface with an anticanonical cycle. *Algebra Number Theory*, 7(6):1481–1504, 2013. [7.4](#)
- [GHK15a] Mark Gross, Paul Hacking, and Sean Keel. Mirror symmetry for log Calabi-Yau surfaces I. *Publ. Math. Inst. Hautes Études Sci.*, 122:65–168, 2015. [1.1](#), [2.2](#), [3.1](#), [3.1](#)
- [GHK15b] Mark Gross, Paul Hacking, and Sean Keel. Moduli of surfaces with an anti-canonical cycle. *Compos. Math.*, 151(2):265–291, 2015. [1.1](#), [1.1](#), [2.1](#), [2.1](#), [2.2](#), [2.2.1](#), [2.5](#), [2.2.1](#), [2.2.2](#)
- [GHKK18] Mark Gross, Paul Hacking, Sean Keel, and Maxim Kontsevich. Canonical bases for cluster algebras. *J. Amer. Math. Soc.*, 31(2):497–608, 2018. [1.1](#)
- [GPS] Sheel Ganatra, John Pardon, and Vivek Shende. Sectorial descent for wrapped Fukaya categories. arXiv:1809.03427v2. [7.3](#)
- [GPS20] Sheel Ganatra, John Pardon, and Vivek Shende. Covariantly functorial wrapped Floer theory on Liouville sectors. *Publ. Math. Inst. Hautes Études Sci.*, 131:73–200, 2020. [5.2](#)
- [Gro85] M. Gromov. Pseudo holomorphic curves in symplectic manifolds. *Invent. Math.*, 82(2):307–347, 1985. [1.3](#), [6.2.2](#), [7.5](#)
- [Gro98] Mark Gross. Special Lagrangian fibrations. I. Topology. In *Integrable systems and algebraic geometry (Kobe/Kyoto, 1997)*, pages 156–193. World Sci. Publ., River Edge, NJ, 1998. [1.2](#)
- [GSW] Honghao Gao, Linhui Shen, and Daping Weng. Augmentations, Fillings, and Clusters. arXiv:2008.10793. [6.2.2](#)
- [Han19] Andrew Hanlon. Monodromy of monomially admissible Fukaya-Seidel categories mirror to toric varieties. *Adv. Math.*, 350:662–746, 2019. [1.3](#), [4.6](#), [5.6](#)
- [Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52. [4.3](#)
- [HH] Andrew Hanlon and Jeffrey Hicks. Functoriality and homological mirror symmetry for toric varieties. arXiv:2010.08817. [1.2](#), [1.3](#)
- [Hic21] Jeffrey Hicks. Tropical Lagrangians in toric del-Pezzo surfaces. *Selecta Math. (N.S.)*, 27(1):Paper No. 3, 50, 2021. [7.8](#)
- [HK] Paul Hacking and Ailsa Keating. Homological mirror symmetry for log Calabi-Yau surfaces. arXiv:2005.05010. [1.1](#), [1.1](#), [1.2](#), [1.3](#), [1.5](#), [1.5](#), [2.8](#), [2.2.2](#), [3.1](#), [3.1](#), [3.3](#), [3.1](#), [3.4](#), [3.2](#), [3.5](#), [3.2.2](#), [3.2.2](#), [3.2.2](#), [4.3](#), [5.1](#), [5.1](#), [6.5](#), [6.2.2](#)
- [HKW] Paul Hacking, Ailsa Keating, and Abigail Ward. in preparation. [1.3](#)
- [Huy06] D. Huybrechts. *Fourier-Mukai transforms in algebraic geometry*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, Oxford, 2006. [1.2](#), [2.3.3](#)
- [Huy16] Daniel Huybrechts. *Lectures on K3 surfaces*, volume 158 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2016. [7.4](#)
- [HW] Andrew Hanlon and Abigail Ward. Tropical Lagrangian sections and Looijenga pairs. in preparation. [1.3](#), [4.6](#), [4.10](#), [5.7](#), [6.2.1](#)
- [IU05] Akira Ishii and Hokuto Uehara. Autoequivalences of derived categories on the minimal resolutions of A_n -singularities on surfaces. *J. Differential Geom.*, 71(3):385–435, 2005. [1.4](#), [1.4](#), [2.1](#), [2.3.3](#)

- [Kea] Ailsa Keating. Families of monotone Lagrangians in Brieskorn–Pham hypersurfaces. arXiv:2002.00286. [6.1](#)
- [Kea15] Ailsa Keating. Lagrangian tori in four-dimensional Milnor fibres. *Geom. Funct. Anal.*, 25(6):1822–1901, 2015. [7.3](#)
- [Kea18] Ailsa Keating. Homological mirror symmetry for hypersurface cusp singularities. *Selecta Math. (N.S.)*, 24(2):1411–1452, 2018. [1.1](#)
- [Kon] M. Kontsevich. Lectures at ENS Paris, Spring 1998. Set of notes taken by J. Bellaïche, J.-F. Dat, I. Marin, G. Racinet and H. Randriambololona. [5.6](#)
- [KS02] Mikhail Khovanov and Paul Seidel. Quivers, Floer cohomology, and braid group actions. *J. Amer. Math. Soc.*, 15(1):203–271, 2002. [1.1](#)
- [KS06] Maxim Kontsevich and Yan Soibelman. Affine structures and non-Archimedean analytic spaces. In *The unity of mathematics*, volume 244 of *Progr. Math.*, pages 321–385. Birkhäuser Boston, Boston, MA, 2006. [1.2](#)
- [KSC04] János Kollár, Karen E. Smith, and Alessio Corti. *Rational and nearly rational varieties*, volume 92 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2004. [1.2](#), [1.5](#), [3.2.1](#)
- [Li] Jennifer Li. A Cone Conjecture for Log Calabi-Yau Surfaces. Doctoral Dissertation, University of Massachusetts, Amherst, 2021. [7.3](#), [7.2](#), [7.2.1](#)
- [Loo81] Eduard Looijenga. Rational surfaces with an anticanonical cycle. *Ann. of Math. (2)*, 114(2):267–322, 1981. [1.1](#), [1.5](#), [2.1](#), [7.2](#)
- [LP04] François Lalonde and Martin Pinsonnault. The topology of the space of symplectic balls in rational 4-manifolds. *Duke Math. J.*, 122(2):347–397, 2004. [7.5](#)
- [LP17] YankıLekili and Alexander Polishchuk. Arithmetic mirror symmetry for genus 1 curves with n marked points. *Selecta Math. (N.S.)*, 23(3):1851–1907, 2017. [1](#)
- [LW12] Tian-Jun Li and Weiwei Wu. Lagrangian spheres, symplectic surfaces and the symplectic mapping class group. *Geom. Topol.*, 16(2):1121–1169, 2012. [1.1](#)
- [McM07] Curtis T. McMullen. Dynamics on blowups of the projective plane. *Publ. Math. Inst. Hautes Études Sci.*, (105):49–89, 2007. [7.4](#)
- [MW18] Cheuk Yu Mak and Weiwei Wu. Dehn twist exact sequences through Lagrangian cobordism. *Compos. Math.*, 154(12):2485–2533, 2018. [7.2](#)
- [Pan17] Yu Pan. Exact Lagrangian fillings of Legendrian $(2, n)$ torus links. *Pacific J. Math.*, 289(2):417–441, 2017. [6.2.2](#), [6.8](#)
- [Per] Timothy Perutz. A symplectic Gysin sequence. arXiv:0807.1863. [1.1](#), [7.2](#)
- [PT20] James Pascaleff and Dmitry Tonkonog. The wall-crossing formula and Lagrangian mutations. *Adv. Math.*, 361:106850, 67, 2020. [1.2](#), [3.2.2](#), [3.2.2](#)
- [Sei] Paul Seidel. Lecture notes on Categorical Dynamics and Symplectic Topology. Available at math.mit.edu/~seidel. [6.4](#)
- [Sei99] Paul Seidel. Lagrangian two-spheres can be symplectically knotted. *J. Differential Geom.*, 52(1):145–171, 1999. [5.1](#)
- [Sei00] Paul Seidel. Graded Lagrangian submanifolds. *Bull. Soc. Math. France*, 128(1):103–149, 2000. [1.1](#), [4.2](#), [7.2](#)
- [Sei01] Paul Seidel. More about vanishing cycles and mutation. In *Symplectic geometry and mirror symmetry (Seoul, 2000)*, pages 429–465. World Sci. Publ., River Edge, NJ, 2001. [6.4](#)
- [Sei03] Paul Seidel. A long exact sequence for symplectic Floer cohomology. *Topology*, 42(5):1003–1063, 2003. [1.1](#), [6.4](#)
- [Sei08] Paul Seidel. *Fukaya categories and Picard-Lefschetz theory*. Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2008. [1.1](#), [5.1](#)

- [Sei17] Paul Seidel. Fukaya A_∞ -structures associated to Lefschetz fibrations. II. In *Algebra, geometry, and physics in the 21st century*, volume 324 of *Progr. Math.*, pages 295–364. Birkhäuser/Springer, Cham, 2017. [5.6](#)
- [Shi90] Tetsuji Shioda. On the Mordell-Weil lattices. *Comment. Math. Univ. St. Paul.*, 39(2):211–240, 1990. [2.2.2](#)
- [Sim] Angelica Simonetti. Equivariant smoothings of cusp singularities. Doctoral Dissertation, University of Massachusetts, Amherst, 2021. [7.2](#)
- [SS20a] Nick Sheridan and Ivan Smith. Symplectic topology of $K3$ surfaces via mirror symmetry. *J. Amer. Math. Soc.*, 33(3):875–915, 2020. [1.1](#), [1.5](#), [7.7](#)
- [SS20b] Vsevolod Shevchishin and Gleb Smirnov. Elliptic diffeomorphisms of symplectic 4-manifolds. *J. Symplectic Geom.*, 18(5):1247–1283, 2020. [1.1](#)
- [ST01] Paul Seidel and Richard Thomas. Braid group actions on derived categories of coherent sheaves. *Duke Math. J.*, 108(1):37–108, 2001. [1.1](#), [2.1](#)
- [STW] Vivek Shende, David Treumann, and Harold Williams. On the combinatorics of exact Lagrangian surfaces. arXiv:1603.07449. [1.1](#), [1.2](#), [3.2.2](#), [6.2.2](#)
- [STWZ19] Vivek Shende, David Treumann, Harold Williams, and Eric Zaslow. Cluster varieties from Legendrian knots. *Duke Math. J.*, 168(15):2801–2871, 2019. [6.2.2](#)
- [Sub10] Aleksandar Subotic. *A monoidal structure for the Fukaya category*. ProQuest LLC, Ann Arbor, MI, 2010. Thesis (Ph.D.)—Harvard University. [1.2](#)
- [Syl19] Zachary Sylvan. On partially wrapped Fukaya categories. *J. Topol.*, 12(2):372–441, 2019. [7.3](#)
- [Sym03] Margaret Symington. Four dimensions from two in symplectic topology. In *Topology and geometry of manifolds (Athens, GA, 2001)*, volume 71 of *Proc. Sympos. Pure Math.*, pages 153–208. Amer. Math. Soc., Providence, RI, 2003. [1.2](#), [3.1](#), [3.4](#), [3.2.2](#), [4](#), [5.1](#), [7.4](#)
- [Tor] Brunella Charlotte Torricelli. Model projective twists and generalised lantern relations. arXiv:2008.02758. [6.1](#)
- [TY90] G. Tian and Shing-Tung Yau. Complete Kähler manifolds with zero Ricci curvature. I. *J. Amer. Math. Soc.*, 3(3):579–609, 1990. [7.4](#)
- [TZ18] David Treumann and Eric Zaslow. Cubic planar graphs and Legendrian surface theory. *Adv. Theor. Math. Phys.*, 22(5):1289–1345, 2018. [6.2.2](#), [6.8](#)
- [Ueh16] Hokuto Uehara. Autoequivalences of derived categories of elliptic surfaces with non-zero Kodaira dimension. *Algebr. Geom.*, 3(5):543–577, 2016. [1.3](#), [1.4](#), [1.4](#), [2.1](#), [2.3.2](#), [2.12](#), [2.3.2](#)
- [Ueh19] Hokuto Uehara. A trichotomy for the autoequivalence groups on smooth projective surfaces. *Trans. Amer. Math. Soc.*, 371(5):3529–3547, 2019. [1.3](#), [1.6](#), [1.4](#), [2.1](#), [2.9](#), [2.3.1](#), [2.10](#), [2.3.3](#)
- [Via14] Renato Vianna. On exotic Lagrangian tori in $\mathbb{C}\mathbb{P}^2$. *Geom. Topol.*, 18(4):2419–2476, 2014. [1.2](#), [3.2.2](#), [3.2.2](#)
- [Via17] Renato Vianna. Infinitely many monotone Lagrangian tori in del Pezzo surfaces. *Selecta Math. (N.S.)*, 23(3):1955–1996, 2017. [1.2](#)
- [WW16] Katrin Wehrheim and Chris T. Woodward. Exact triangle for fibered Dehn twists. *Res. Math. Sci.*, 3:Paper No. 17, 75, 2016. [1.1](#), [7.2](#)

DEPARTMENT OF MATHEMATICS AND STATISTICS, LEDERLE GRADUATE RESEARCH TOWER, UNIVERSITY OF MASSACHUSETTS, AMHERST, MA 01003-9305

Email address: `hacking@math.umass.edu`

DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS, CENTRE FOR MATHEMATICAL SCIENCES, UNIVERSITY OF CAMBRIDGE, WILBERFORCE ROAD, CAMBRIDGE, CB3 0WB

Email address: `amk50@cam.ac.uk`