

DEPARTMENT OF MATHEMATICS AND STATISTICS
UNIVERSITY OF MASSACHUSETTS

MATH 233

MAKE-UP & DS EXAM 2

Fall 2018

NAME: _____ Student ID: _____

Section Number: _____ Instructor's Name: _____

In this exam there are NINE PAGES, including this one, and SEVEN problems. Make sure you have them all. You must SHOW ALL WORK. Calculators, textbooks, class notes, formula sheets, etc. are NOT allowed. BOX your final answers.

- | | | |
|-------|-------|-------|
| 1. | (14) | _____ |
| 2. | (14) | _____ |
| 3. | (14) | _____ |
| 4. | (14) | _____ |
| 5. | (14) | _____ |
| 6. | (15) | _____ |
| 7. | (15) | _____ |
| Total | (100) | _____ |

1. A particle moves along the curve

$$\mathbf{r}(t) = (t^4/4, (\sqrt{6}/3)t^3, (3/2)t^2).$$

Find the length of the path traveled by the particle between $t = 1$ and $t = 2$.

2. Find the position vector function of a particle that has an acceleration function

$$\mathbf{a}(t) = \cos(t/2)\mathbf{i} + \mathbf{k},$$

an initial velocity $\mathbf{v}(0) = 3\mathbf{j}$, and an initial position $\mathbf{r}(0) = \mathbf{0}$.

3. Consider the function

$$f(x, y) = \frac{x^2y}{x^4 + 2y^2}.$$

- (a) Find the limit of the $f(x, y)$ as (x, y) approaches the origin *along a straight line* of slope m , where m is a real number.
- (b) Find the limit of the $f(x, y)$ as (x, y) approaches the origin *along the curve* $y = x^2$.
- (c) Does $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$ exist? Explain why or why not.

4. (a) Explain why $f(x, y) = \sqrt{x^2 + y^2 + 1}$ is differentiable at $(4, 8)$, and then find the equation of the tangent plane to graph of $z = f(x, y)$ when $x = 4$ and $y = 8$.

(b) The volume of a square pyramid is measured as 30 cubic centimeters with a possible error of ± 1 cubic centimeters. Its height is measured as 10 centimeters, with a possible error of ± 0.3 centimeters. Use differentials to estimate the maximum error in calculating the side length of the square base from the measured volume and height. You must include units in your final answer. (Recall that the volume of a pyramid with a square base of side length l and height h is $V = \frac{1}{3}l^2h$.)

5. (a) Find the directional derivative of the function $f(x, y) = xe^{-y}$ in the direction of $\mathbf{v} = \langle 1, 2 \rangle$ at the point $P = (1, 0)$.
- (b) Find the maximum possible directional derivative of $f(x, y) = xe^{-y}$ at the point $P = (1, 0)$ and the unit vector in the direction in which it occurs.

6. Find and classify (i.e., local maximum, local minimum, saddle point, or inconclusive) all critical points of the function:

$$f(x, y) = y^3 - \frac{3}{2}y^2 + \frac{3}{2}x^2 - 3xy + 5.$$

7. Use the method of Lagrange multipliers to find the absolute maximum and absolute minimum values of the function $f(x, y) = e^{xy}$ on the curve given by $x^2 + y^2 = 2$.

Scratch paper