

Conformal Structures and Necksizes of Embedded Constant Mean Curvature Surfaces

Rob Kusner

Dedicated to my father and mentor, David B. Kusner, on the occasion of his seventieth birthday

ABSTRACT. Let $\mathcal{M} = \mathcal{M}_{g,k}$ denote the space of properly embedded (or Alexandrov immersed) constant mean curvature (CMC) surfaces of genus g with k (labeled) ends, modulo rigid motions, endowed with the real analytic structure described in [20]. Let $\mathcal{P} = \mathcal{P}_{g,k} = \mathcal{R}_{g,k} \times \mathbb{R}_+^k$ be the space of parabolic structures over Riemann surfaces of genus g with k (marked) punctures, the real analytic structure coming from the $3g - 3 + k$ local complex analytic coordinates on the Riemann moduli space $\mathcal{R}_{g,k}$. Then the *parabolic classifying map*, $\Phi : \mathcal{M} \rightarrow \mathcal{P}$, which assigns to a CMC surface its induced conformal structure and asymptotic necksizes, is a proper, real analytic map. It follows that Φ is closed and in particular has closed image. For genus $g = 0$, this can be used to show that every conformal type of multiply punctured Riemann sphere occurs as a CMC surface, and—under a nondegeneracy hypothesis—that Φ has a well-defined (mod 2) degree. This degree vanishes, so generically an *even* number of CMC surfaces realize any given conformal structure and asymptotic necksizes (compare [11, 12] for the case $k = 3$).

Introduction

Besides their beauty and variety, perhaps the most important reason minimal surfaces have been so thoroughly investigated is because they can be conformally parametrized via holomorphic functions of a complex variable, a long-familiar tool to many mathematicians. This representation theory was worked out by Enneper and Weierstrass in the middle of the 19th century, leading to important local results and interesting periodic examples. But not until the 1960s did Osserman [27] prepare the way for a global theory by showing that a complete minimal surface with finite total curvature is conformally parametrized by meromorphic data on a finitely punctured, finite genus Riemann surface. The global Enneper–Weierstrass representation enables one to employ algebro-geometric methods from the theory of complex curves to study such minimal surfaces. Over the past two decades it has been used in conjunction with analytic methods by a number of authors—many represented in this volume—to prove deep and striking results about properly embedded minimal surfaces of finite total curvature or finite topology (finite genus and a finite number of ends).

The theory of complete, properly embedded surfaces with *nonzero* constant mean curvature (CMC) has developed more recently and without the benefit of these holomorphic methods. While more than a century elapsed between Delaunay's classification [7] of the CMC rotation surfaces (two ends, genus zero) and Alexandrov's [2] proof that the only compact (zero ends) CMC surface is the round sphere, only in the past decade have interesting noncompact examples of finite topology CMC surfaces (with three or more ends) been constructed by Kapouleas [16], using difficult analytic methods. At about the same time, the first steps toward a global theory of these surfaces were taken by Meeks and by Korevaar, Kusner and Solomon: they found obstructions to the existence of CMC surfaces with finite topology—none have one end [24], two-ended surfaces are Delaunay unduloids [19]—and developed an asymptotic theory for these surfaces. In fact the fundamental result of [19] is that each (annular) end of a CMC surface has an unduloid asymptote. An immediate consequence is that a finite topology CMC surface can be conformally parametrized by a finitely punctured Riemann surface of finite genus.




FIGURE 1. Equilateral triunduloid and tetrunduloid (images by Nick Schmitt, GANG).

Can such a global conformal parametrization be made explicit—enough to prove something new about CMC surfaces, or at least to experiment with them using a computer? Using a loop-group analogue of the Enneper–Weierstrass representation—dubbed the DPW representation after its discoverers, Dorfmeister, Pedit and Wu [8]—Schmitt has recently taken up this question at GANG in Amherst (see Figure 1, and www.gang.umass.edu). He adapted the DPW representation to construct, in terms of explicit meromorphic data on a punctured Riemann surface, conformal immersions of finite topology constant mean curvature surfaces whose ends are asymptotically Delaunay unduloids (or nodoids) [17, 30]. This approach is still under development, and little is known about how to prove that the examples it constructs have the symmetries or embeddedness properties required. Nevertheless, it appears to recapture the earlier results of Große-Brauckmann, Kusner and Sullivan [11, 12] about three-ended CMC surfaces of genus zero, the *triunduloids*.

Our investigations [11, 12] made use of a special classifying map from the space of all triunduloids to an open three-ball. We relied heavily on the real analytic variety structure of CMC moduli space developed by Kusner, Mazzeo, and Pollack [20],

as well as the compactness results of Korevaar and Kusner [18], to prove this map is a homeomorphism. The present paper also uses these transcendental methods, and was motivated by a desire to weave these together with the holomorphic methods arising from the DPW representation in order to better comprehend the fabric of CMC moduli spaces.

The main result of this paper is a properness theorem for a general classifying map which assigns a CMC surface its induced conformal structure and asymptotic necksizes. As noted in the abstract, consequences of this theorem include information about which conformal structures and necksizes—input data for the DPW representation—can arise for CMC surfaces, as well as about how many surfaces might share given data. In particular, we show that these surfaces generally occur in pairs, a phenomenon first observed by Karcher and Große-Brauckmann [10] for surfaces with special symmetry. Several outstanding open problems about the moduli spaces of CMC surfaces are also discussed.

There is some interplay between our results and those of Mazzeo, Pacard and Pollack, so here an effort is made to focus on what is complementary; the reader is encouraged to refer to their nice paper [23], as well as to [12], for additional background and related material. The author wishes to thank Rafe Mazzeo, Frank Pacard, Dan Pollack, Jesse Ratzkin and Rick Schoen for stimulating discussions, and Karsten Große-Brauckmann for much thoughtful advice. Thanks are also due to Nicos Kapouleas, Hermann Karcher, Nick Korevaar, Remi Langevin, Franz Pedit, Wayne Rossman, Nick Schmitt, John Sullivan and Mike Wolf for their interest and comments, and especially to David Hoffman and Harold Rosenberg for their invitation to present this at the 2001 Clay/MSRI summer workshop. The research reported here was supported in part by NSF grant DMS-0076085.

1. Background on the CMC Moduli Space

Let $\mathcal{M} = \mathcal{M}_{g,k}$ be the moduli space of all CMC surfaces of genus g with k (labeled) ends, modulo rigid motions. Here, as usual, we scale the mean curvature to be 1, and include in \mathcal{M} not only the embedded surfaces, but also those which are immersed in the sense of Alexandrov: these are boundaries of properly immersed 3-dimensional manifolds (compare [3, 12, 19]). Some of the results mentioned in the introduction can then be expressed in terms of \mathcal{M} as follows:

- $\mathcal{M}_{0,0}$ is a point, represented by any unit sphere, and $\mathcal{M}_{g,0}$ is empty for all $g > 0$ [2];
- $\mathcal{M}_{g,1}$ is empty for every g [24];
- $\mathcal{M}_{0,2}$ consists of the unduloids (up to rigid motion), but $\mathcal{M}_{g,2}$ is empty for all $g > 0$ [19];
- $\mathcal{M}_{g,k}$ is nonempty for every $g \geq 0$, $k > 2$ [16]—for example, $\mathcal{M}_{0,3}$ is a 3-ball [11, 12];
- $\mathcal{M}_{g,k}$ is a real analytic variety of (virtual) dimension $3k - 6$, for $k > 2$ [20].

To understand this real analytic structure on \mathcal{M} , it will be useful to introduce the *pre-moduli* space $\widetilde{\mathcal{M}}$, that is, the space of CMC surfaces *before* dividing by rigid motions. We also let $\mathcal{U} = \widetilde{\mathcal{M}}_{0,2}$ denote the pre-moduli space of all unduloids. The discussion below will describe the real analytic variety structure on $\widetilde{\mathcal{M}}$ in terms

of an explicit real analytic manifold structure on \mathcal{U} , in order to demonstrate the following (almost tautological) result:

PROPOSITION 1.1. *The map $\widetilde{\mathcal{M}}_{g,k} \rightarrow \mathcal{U}^k$ which assigns a CMC surface its k unduloid asymptotes is real analytic, and is equivariant with respect to the actions of rigid motions. Moreover, there is a real analytic necksize function $s : \mathcal{U} \rightarrow \mathbb{R}_+$, taking values in $(0, 1]$. It is invariant under rigid motions, and so gives rise to a real analytic asymptotic necksize map $(s_1, \dots, s_k) : \mathcal{M}_{g,k} \rightarrow \mathbb{R}_+^k$, taking values in $(0, 1]^k$.*

PROOF. The equivariance is clear when we take the diagonal action by rigid motions on the k -fold product \mathcal{U}^k : as a CMC surface undergoes a rigid motion, each unduloid asymptote undergoes the same motion. Furthermore, once s is defined, it should be clear that (s_1, \dots, s_k) is defined by composition with the asymptotes map and passage to the quotient by rigid motions.

Recall that an unduloid [19] is a periodic CMC surface of revolution determined by its axis, its neckphase—where along the axis it is nearest the axis—and its necksize. (Of course, modulo rigid motions, an unduloid is specified by its necksize alone.) So the space \mathcal{U} is naturally a fiber bundle over the space $T\mathbb{S}^2$ of (oriented) lines in \mathbb{R}^3 . The base space $T\mathbb{S}^2$ has a natural analytic structure as a homogeneous space for the group of rigid motions. The total space \mathcal{U} will then be expressed as a 6-dimensional real analytic manifold provided an analytic manifold structure can be exhibited on the typical fiber \mathcal{D} , consisting of all unduloids with a fixed, oriented axis (the two ends of the unduloid are labeled according to the axis orientation).

Observe that \mathcal{D} is topologically an open disk, with necksize and neckphase as polar coordinates. The cylinder, whose necksize is maximal and whose neckphase is not well defined, is at the center of \mathcal{D} , fixed under rotation (neckphase-shift). The boundary of \mathcal{D} corresponds to the zero necksize limits of unduloids—chains of spheres along the given axis. The natural analytic structure on \mathcal{D} comes from its (locally homogeneous) realization as an open subset of the unit sphere \mathbb{S}^2 , indeed as $\mathbb{S}^2 - \{p\}$ for a single point p .

This spherical picture of \mathcal{D} also played a key role in [12], leading to a natural interpretation of necksize—there referring to the length of the shortest nontrivial loop on an unduloid—as the spherical distance n between p and another point $q \in \mathcal{D} = \mathbb{S}^2 - \{p\}$. Unfortunately, n is not an analytic coordinate on \mathcal{D} : at the cylinder (corresponding to q antipodal to p) the function n is only Lipschitz. This will be remedied by reinterpreting *necksize* here to mean $s = \frac{1}{2}(1 - \cos n)$.

Although s may seem geometrically less natural than n , it turns out to be more natural in other ways. For example, being a height function on \mathbb{S}^2 in the direction p , it is symplectically dual to neckphase (angle around p) for the homogeneous symplectic structure on \mathbb{S}^2 . For the present paper, however, the important point is that s is analytic with respect to the natural (locally homogeneous) analytic structure on \mathcal{U} , and has the invariance properties and range indicated in the statement of the proposition. In fact, we could have equally well used the *weight* or *force* modulus [19] of an unduloid, $f = n(2\pi - n)$, which is also an analytic coordinate on \mathcal{U} , even at the cylinder ($n = \pi$).

The tangent space to \mathcal{U} at an unduloid can be identified with the 6-dimensional linear space \mathcal{V} of *geometric* Jacobi fields (see [19, 20]). Thus \mathcal{V} gives analytic local coordinates on \mathcal{U} in a neighborhood of this unduloid. We shall pretend that \mathcal{V} defines global analytic coordinates on \mathcal{U} in what follows, but since checking

analyticity is a local condition, this pretense only amounts to a mild abuse of notation.

Using the asymptotics result [19], one can construct [20] a $6k$ -dimensional linear space \mathcal{W} of functions on any surface $\Sigma \in \widetilde{\mathcal{M}}_{g,k}$ which grow at each end of Σ like those in \mathcal{V} . There are various ways to realize such a \mathcal{W} — either as an orthogonal complement (in an appropriate Hilbert space, compare [20]) to the functions which decay on every end of Σ , or as a quotient space (compare [22]) by such functions — but the salient point is that there is a linear isomorphism from \mathcal{W} to \mathcal{V}^k , and thus a real analytic map from \mathcal{W} to \mathcal{U}^k .

So in order to establish the proposition above, it would suffice to analytically embed $\widetilde{\mathcal{M}}$ into \mathcal{W} , at least locally, in the natural way: at a CMC surface Σ , represent any nearby CMC surface in $\widetilde{\mathcal{M}}$ via its asymptotic behavior, which is encoded by a point of \mathcal{W} . One of the main results in [20] is that this is possible when Σ is *nondegenerate*, that is, provided Σ supports no nontrivial L^2 -Jacobi fields. In this case all nearby CMC surfaces form an analytic manifold of dimension $3k$ — half that of \mathcal{W} — and $\widetilde{\mathcal{M}}_{g,k}$ is (locally) a Lagrangian submanifold with respect to a natural symplectic structure defined [20] on \mathcal{W} (or on \mathcal{V}^k , or even directly on \mathcal{U}^k).

For the general case, a neighborhood of Σ in $\widetilde{\mathcal{M}}$ may no longer analytically embed in \mathcal{W} , but it can still be embedded as a variety in a larger linear space $\mathcal{W} \times \mathcal{K}$ where \mathcal{K} is a finite-dimensional linear space accounting for the L^2 -Jacobi fields on Σ (compare [20]). Clearly the composition of analytic maps — inclusion of this variety in $\mathcal{W} \times \mathcal{K}$, followed by projection to \mathcal{W} , then the linear isomorphism to \mathcal{V}^k , and finally, the analytic local coordinates map into \mathcal{U}^k — is the asymptotes map, which is thus analytic. □

2. A Proper Classifying Map and Its Consequences

One can investigate the topology of \mathcal{M} by studying maps with special properties from \mathcal{M} to other known spaces. As noted in the introduction, this is the method used to classify the triunduloids: there is a homeomorphism Ψ from $\mathcal{M}_{0,3}$ to an open 3-ball, realized by the space of ordered triples of points on \mathbb{S}^2 (up to rotations): Ψ identifies the asymptotic necksizes of the triunduloid with the spherical distances for the triple [11, 12]. Unfortunately, the triunduloid classifying map Ψ requires the existence of a mirror symmetry which is not necessarily present for CMC surfaces with four or more ends, and so we will introduce a more general classifying map here to investigate them.

We have already observed that the *necksize map* $(s_1, \dots, s_k) : \mathcal{M}_{g,k} \rightarrow \mathbb{R}_+^k$, is real analytic. Now consider the *forgetful map* $\phi : \mathcal{M}_{g,k} \rightarrow \mathcal{R}_{g,k}$ which assigns the CMC surface Σ the conformal class $[\Sigma]$ of its induced metric from \mathbb{R}^3 . In other words, ϕ “forgets” everything but the underlying punctured Riemann surface. It is well known (compare [1, 15, 32, 31]) that the Riemann moduli space $\mathcal{R}_{g,k}$ can be given a real analytic structure in two (equivalent) ways, either in terms of holomorphic quadratic differentials with prescribed poles at the punctures ($3g - 3 + k$ complex analytic local coordinates coming from the cotangent bundle to $\mathcal{R}_{g,k}$) or in terms of finite-area hyperbolic metrics ($6g - 6 + 2k$ real analytic Fenchel–Nielsen coordinates). One could exploit the Hopf differential for the Gauss map of a constant mean curvature surface to derive the following result via holomorphic quadratic differentials, but fortunately Mazzeo, Pacard and Pollack have already derived this [23] by proving a nontrivial result in semilinear elliptic partial differential equations;

they show the unique hyperbolic metric on Σ —conformal to its induced metric as a CMC surface—depends analytically on Σ , and thus [23]:

PROPOSITION 2.1. *The forgetful map ϕ is real analytic.*

Although it is tempting to work with the forgetful map ϕ alone, there are at least two reasons why another map will be more suitable. First, in order to find homotopy invariants for maps between noncompact spaces, the maps are generally required to be proper. Unfortunately ϕ itself is not proper: compare $\mathcal{R}_{0,3}$, which is a single point (and thus compact), to its preimage $\mathcal{M}_{0,3}$ under ϕ , which is an open (and thus noncompact) 3-ball. In general ϕ fails to be proper because a sequence of CMC surfaces diverges in \mathcal{M} by having asymptotic necksizes tend to zero, while remaining in a compact family of conformal types. Second, the meromorphic data for the DPW representation of a CMC surface with asymptotically unduloidal ends can be geometrically interpreted in terms of flat connections on a rank-2 complex vector bundle over a punctured Riemann surface, with parabolic holonomy determined by a positive number (related to the asymptotic necksize) at each of the punctures (compare [5, 30]).

These considerations suggest extending the Riemann moduli space to the *parabolic moduli space* $\mathcal{P}_{g,k} = \mathcal{R}_{g,k} \times \mathbb{R}_+^k$ which assigns a positive real number to each puncture of $[\Sigma]$, and defining the *parabolic classifying map*

$$\Phi : \mathcal{M}_{g,k} \rightarrow \mathcal{P}_{g,k} = \mathcal{R}_{g,k} \times \mathbb{R}_+^k$$

as the product of the forgetful map ϕ with the asymptotic necksize map (s_1, \dots, s_k) . The main result of this paper is the following:

THEOREM 2.2. *The parabolic classifying map $\Phi : \mathcal{M} \rightarrow \mathcal{P}$ is real analytic and proper.*

Note that the real analyticity of Φ follows immediately from Propositions 1.1 and 2.1. Before giving a proof of properness (see Section 3), we first point out several consequences:

COROLLARY 2.3. *The parabolic classifying map $\Phi : \mathcal{M} \rightarrow \mathcal{P}$ is closed. In fact, its image is a closed real analytic subvariety, which represents a (mod 2) homology class in \mathcal{P} . The forgetful map ϕ also has closed image in \mathcal{R} .*

PROOF. Since \mathcal{P} is a compactly generated Hausdorff space, we can apply a lemma from general topology (see [12], Section 5) to show that any proper map to \mathcal{P} is closed; in particular, Φ has closed image $\Phi(\mathcal{M}) \subset \mathcal{P}$. The homology class in \mathcal{P} is carried by the image of Φ , or more precisely, by the proper (mod 2) cycle which is the push-forward via Φ of the fundamental (mod 2) cycle of the real analytic variety \mathcal{M} (compare [6, 33]).

The forgetful map ϕ is the composition of Φ with the projection from \mathcal{P} to \mathcal{R} . Unfortunately, projection is generally not a closed map, so the proof that the image of ϕ is closed does not follow immediately from the fact that the image of Φ is closed. Nevertheless, the analyticity of the forgetful map ϕ (Proposition 2.1) suffices to show it has closed image (compare [23]). \square

Next, observe that any component of $\mathcal{M}_{g,k}$ containing a nondegenerate CMC surface is a real analytic variety of dimension $3k - 6$ [20], and this coincides with $6g - 6 + 3k$ —the dimension of $\mathcal{P}_{g,k}$ —precisely when $g = 0$. This is a third important

reason to introduce \mathcal{P} , since it leads to a degree theory, at least in case of genus zero surfaces (we shall assume $g = 0$ for the remainder of this section, with the exception of the final remark):

THEOREM 2.4. *For any component \mathcal{N} of $\mathcal{M}_{0,k}$ containing a nondegenerate CMC surface, the restriction of Φ to \mathcal{N} has a well-defined degree (mod 2). Moreover, this degree is zero.*

PROOF. Because the domain \mathcal{N} and the range \mathcal{P} have the same dimension, the ideas in [6, 12, 33] let us define this degree as the number of pre-images (mod 2) for any regular value of Φ . The connectedness of \mathcal{P} and properness of Φ imply that the pre-images of any two regular values can be joined by a compact, 1-dimensional (semi-analytic) variety in \mathcal{N} , which necessarily has an even number of endpoints, so the numbers of pre-images agree (mod 2). Then the obvious upper bound on necksize by that of a cylinder shows that this (mod 2)-degree must vanish. \square

In particular, it follows that genus zero CMC surfaces generically occur in pairs:

COROLLARY 2.5. *For any nondegenerate surface $\Sigma \in \mathcal{M}_{0,k}$ which is a regular point of Φ , there is a corresponding surface $\Sigma' \in \mathcal{M}_{0,k}$ with the same conformal structure and necksizes.*

Of course, there is no reason to expect there are exactly *two* such surfaces with the same data, only an *even* number (see Section 5 for further discussion of this phenomenon).

Finally, we observe that every conformal type of punctured Riemann sphere is realized by a CMC surface. Although we make no use of this below, this observation is useful from the perspective of the DPW representation, since it guarantees the search for DPW data on a particular punctured Riemann sphere will (in principle) always succeed. Although this is also proven via a rather different argument—using end attachment at (almost) any point of a (nondegenerate) CMC surface, and induction on the number of ends—in [23], it is interesting to present this direct argument, based on ideas coming from Kapouleas’ original construction of CMC surfaces [16]:

THEOREM 2.6. *The forgetful map $\phi : \mathcal{M}_{0,k} \rightarrow \mathcal{R}_{0,k}$ is surjective.*

PROOF. From the second part of Corollary 2.3, it suffices to show that a dense set of conformal types is realized. Using a little linear algebra, it is not hard to see that, up to conformal transformations of \mathbb{S}^2 , any configuration of $k > 1$ points ξ_1, \dots, ξ_k on \mathbb{S}^2 can be balanced with *positive* weights: that is, regarding the ξ_j as unit vectors in \mathbb{R}^3 , there exists a positive solution f_1, \dots, f_k to the linear relation (force balancing)

$$f_1 \xi_1 + \dots + f_k \xi_k = 0.$$

For sufficiently small forces f_j , the Kapouleas construction [16] then gives a CMC surface Σ of genus zero with k ends asymptotic to unduloids whose j th axis is approximately in the direction ξ_j and whose corresponding necksize is approximately f_j . Because Σ is obtained by quasiconformally attaching punctured disks (corresponding to the unduloid ends) at the boundaries of arbitrarily small disks about each ξ_j , its conformal structure $[\Sigma]$ can be made to lie within any prescribed neighborhood of $[\mathbb{S}^2 - \{\xi_1, \dots, \xi_k\}]$ in $\mathcal{R}_{0,k}$. \square

It is quite easy to see that ϕ cannot be surjective for $g > 0$. For instance, when $k = 3$ a CMC surface Σ must have a plane of reflection symmetry, so the Riemann surface $[\Sigma] = \phi(\Sigma)$ must have a certain anti-conformal involution with more than g component curves in the fixed point set; this necessary condition, for example, allows only rectangular tori in $\mathcal{M}_{1,3}$.

There is an interesting generalization of Φ and $\mathcal{M}_{g,k}$ which admits results analogous to Theorems 2.2 and 2.4 (and their Corollaries) in case $g > 0$. To understand this, recall that any $\Sigma \in \mathcal{M}_{g,k}$ bounds an immersed genus- g handlebody Ω ; because Σ closes when we follow a loop in Ω means we have the *trivial* representation of $\pi_1(\Omega)$ into the group of rigid motions. The new idea is to consider *all* such representations, where Σ may no longer have closed periods around the g generators of $\pi_1(\Omega)$; the corresponding space $\mathcal{M}_{g,k}^*$ of *half-periodic* CMC surfaces forms a variety whose (virtual) dimension is $6g$ more than that of $\mathcal{M}_{g,k}$. A fundamental domain for a half-periodic surface still defines a punctured Riemann surface, and it still has k asymptotic necksizes; thus we get an *extended parabolic classifying map* Φ^* in the obvious way:

REMARK 2.7. *The extended parabolic classifying map $\Phi^* : \mathcal{M}_{g,k}^* \rightarrow \mathcal{P}_{g,k}$ is real analytic and proper. Any component \mathcal{N}^* of \mathcal{M}^* containing a nondegenerate half-periodic CMC surface is $(3k - 6 + 6g)$ -dimensional (the same as for \mathcal{P}), and thus the restriction of Φ^* to \mathcal{N}^* has a well-defined (mod 2) degree, which must vanish.*

We plan to explore the consequences of these observations about Φ^* , along with the problem of closing periods of $\Sigma \in \mathcal{M}_{g,k}^*$, in a future paper.

3. Proof of Properness

The proof that the parabolic classifying map $\Phi : \mathcal{M} \rightarrow \mathcal{P}$ is proper relies on the *a priori* estimates for CMC surfaces developed in [18], namely:

LEMMA 3.1. *Any $\Sigma \in \mathcal{M}_{g,k}$ lies in a uniform tubular neighborhood of a piecewise linear 1-complex with k rays and at most $k + 3g - 3$ finite segments. Moreover, a sequence $\Sigma(i)$ of CMC surfaces, all of which lie in a compact family of uniform tubular neighborhoods, diverges in $\mathcal{M}_{g,k}$ if and only if the length $\ell(i)$ of the shortest nontrivial loop on $\Sigma(i)$ tends to zero (compare [18]).*

A compact family of uniform tubular neighborhoods is characterized by requiring the lengths of all segments to be uniformly bounded above, although these lengths may go to zero, as may the angles between the rays or edges. For sequences of CMC surfaces $\Sigma(i)$ lying in such a compact family of tubular neighborhoods, uniform linear area and total curvature estimates [18] still hold, and uniform pointwise curvature estimates also hold under the hypothesis that the $\ell(i)$ are bounded away from zero (using blow-up arguments of the kind we outline below).

The proof also depends on a description of divergent sequences of punctured Riemann surfaces in terms of the conformal moduli of certain nontrivial annuli. We say that an embedded annulus A on a punctured Riemann surface is *essential* (or nonperipheral) provided it is not homotopic to a single point of the surface, nor to a single puncture. Such an annulus then has a (necessarily finite) conformal modulus $[A] = m \in (1, \infty)$ defined by conformally mapping A into \mathbb{C} so that one boundary component goes to the unit circle, and the other goes to the circle of radius m . The basic fact which we will use is the following:

LEMMA 3.2. *A sequence of punctured Riemann surfaces $[\Sigma(i)]$ diverges in $\mathcal{R}_{g,k}$ if and only if there exist essential annuli $A(i) \subset \Sigma(i)$ whose moduli $[A(i)] = m(i) \rightarrow \infty$ as $i \rightarrow \infty$.*

PROOF. This is a standard result in Teichmüller theory, using extremal length (compare, for instance, [32]) or Fenchel–Nielsen coordinates (compare [1, 15, 31]), although it seems difficult to find the exact statement in the literature, particularly for the case of a punctured Riemann surface. In effect, the existence of a family of essential annuli $A(i) \subset \Sigma(i)$ whose moduli diverges to infinity is equivalent to having a family of geodesics (in the homotopy class of $A(i)$) whose length in the conformal hyperbolic metric on $[\Sigma(i)]$ tends to zero. The Mumford–Bers compactness theorem (compare [26, 4]) implies that only these $[\Sigma(i)]$ diverge in $\mathcal{R}_{g,k}$. \square

Proof of the properness theorem. To show that the classifying map Φ is proper it will suffice to prove the contrapositive, namely, that a divergent sequence $\Sigma(i)$ in $\mathcal{M}_{g,k}$, whose asymptotic necksizes $s_1(i), \dots, s_k(i)$ are uniformly bounded away from zero, must contain divergent essential annuli as above, implying $\Phi(\Sigma(i))$ is divergent in $\mathcal{P}_{g,k}$.

So suppose a sequence $\Sigma(i)$ diverges in \mathcal{M} , with all asymptotic necksizes $s_j(i) \geq \ell > 0$. Then we have two (non-exclusive) alternatives:

- $L(i) \rightarrow \infty$, where $L(i)$ is the length of the longest finite segment in the 1-complex whose uniform tubular neighborhood contains $\Sigma(i)$; or
- $\ell(i) \rightarrow 0$, where $\ell(i)$ is the length of some nontrivial closed curve, which is not homotopic to an end of $\Sigma(i)$.

In the first case, the Alexandrov symmetrization method [18, 19] shows there is a sequence of essential annuli $A(i) \subset \Sigma(i)$ which are approximately unduloidal on greater and greater lengths (comparable to $L(i) \rightarrow \infty$, though the constants are not explicitly computable). Since any unduloid is periodic, the essential annuli approximating an unduloid have moduli $[A(i)]$ which grow exponentially with length, and thus diverge.

In the second case, we make a blow-up argument (compare [12, 18, 19]) to get a non-flat, finite total curvature minimal surface M bounding an immersed 3-manifold. There are several subcases to consider.

First, suppose M is embedded, so its top and bottom ends are catenoidal. Let A be a punctured-disk representative for such an end on M . Thus as $i \rightarrow \infty$, the end A is approximated by rescaled copies of annuli $A(i) \subset \Sigma(i)$ with moduli $[A(i)] \rightarrow [A] = \infty$. Since A is catenoidal, it has nonzero force (see [12, 18, 19]), as must the $A(i)$, which implies they are homologically nontrivial. And although the forces (and thus the necksizes) of the $A(i)$ tend to zero, because we are assuming the necksizes of the ends of $\Sigma(i)$ are bounded away from zero, it follows that the $A(i)$ are not homotopic to any end of $\Sigma(i)$, and so must be essential.

Next, suppose the blow-up minimal surface M is only immersed. If each end of M is simple, then each must be either catenoidal or planar. An application of the strong half-space theorem [14], using the immersed 3-manifold in place of \mathbb{R}^3 , shows that not all ends of M can be planar; so we get a catenoidal end with nonzero force, and a punctured-disk representative A , to use just as in the argument above.

Finally, in case an end of the blow-up minimal surface M is non-simple, we use a “slicing trick” (compare with the instructive example in Section 4): we find a smoothly embedded plane P in \mathbb{R}^3 , which meets a representative annulus A for

this end in a curve which cannot bound a compact immersed surface on P ; it will then follow that A must have been approximated by rescaled essential annuli $A(i)$ on $\Sigma(i)$. Such a P can be constructed by taking a large radius cylinder with axis parallel to the asymptotic unit normal of A , and capping off by a hemisphere very high up on this cylinder (choose this normal to be on the side which bounds the immersed 3-manifold and think of this side as up). Its curve of intersection with the rescaled approximating $A(i)$ has turning number in P equal to the asymptotic winding number of A , which is at least 2. Endow P with an intrinsically flat metric and apply the Gauss–Bonnet formula to see this curve cannot bound a compact immersed surface in P : its winding number would equal the Euler number of a connected surface with boundary, which is at most 1. This shows that this curve of intersection cannot bound in the 3-manifold itself. It follows that the annuli $A(i) \subset \Sigma(i)$ approximating A must have been homotopically nontrivial in the immersed 3-manifold bounded by $\Sigma(i)$ as well, and thus essential in $\Sigma(i)$.

This completes the proof of our main result.

4. An Instructive Example: the Enneper Surface

The reader may find it instructive to apply this slicing trick to decide whether a particular immersed surface can actually be extended to a properly immersed 3-manifold. Take, for example, the Enneper surface, a minimally immersed \mathbb{R}^2 in \mathbb{R}^3 ; it has one end, of winding number 3. Note that, in general, the Frankel–Lawson [9, 21] argument shows the inclusion of the bounding minimal surface into the 3-manifold induces an epimorphism on fundamental groups. So if Enneper were to bound, it would bound an immersed 3-ball (with one boundary puncture, corresponding to the one end of Enneper). But the slicing trick implies that the intersection curve of Enneper with a slicing plane P , constructed as above, has turning number 3, and therefore the curve would be homotopically nontrivial in the 3-ball, which is absurd. Thus:

COROLLARY 4.1. *The Enneper surface is not immersed in the sense of Alexandrov.*

5. Some Open Problems

To conclude, it may be interesting to contemplate the following (open) problems which are related to the results above:

- What is the image of the parabolic classifying map Φ in \mathcal{P} , or its intersection with a slice of fixed conformal type? In other words, how does the set of allowable necksizes depend on the underlying punctured Riemann surface? For $g = 0$ and $k = 3$, there is only one conformal type, so these questions coincide: the image of Φ is known to be the simplex determined by the spherical triangle inequalities for the necksizes of a triunduloid [12]. The general case seems closely related to the Biswas inequalities [5, 30] which arise in the study of flat connections on parabolic vector bundles.
- Is the cardinality of $\Phi^{-1}([\Sigma])$ finite, and is $2^{|\chi(\Sigma)|}$ an upper bound? The absolute value of the Euler characteristic $|\chi(\Sigma)| = 2g - 2 + k$ is the number of trousers in a decomposition of Σ . Thus the possible choices of “innie” or “outie” configurations at each trousers nexus saturates this bound. (This whimsical terminology — suggested by the proximity of each trousers nexus to the umbilic points on the surface —

describes either a neck-like or bubble-like geometry near the trousers nexus.) Again, for $g = 0$ and $k = 3$ we have $|\chi(\Sigma)| = 1$, and this bound is sharp [12]: Φ is two-to-one, except at the maximal necksize “fold” where it is one-to-one.

- Is $\mathcal{M}_{g,k}$ connected? And—at least on some connected component of \mathcal{M} —how far from an isomorphism is the homomorphism on fundamental groups induced by Φ —or, equivalently, by ϕ ? In case $g = 0$ it follows from [12] and [29] (or more directly from [23]) that, on what is presumably the only component—the one containing the *coplanar* CMC surfaces [13]—of $\mathcal{M}_{0,k}$, this is an epimorphism for any k .

- Does $\mathcal{M}_{g,k}$ carry a complete metric of nonpositive curvature, in some suitable sense, and is it thus an Eilenberg–MacLane $K(\pi, 1)$ -space? The identification of $\mathcal{M}_{0,3}$ with the (hyperbolic) 3-ball in [12] is very suggestive, but it is not clear where to find a metric (hyperbolic or otherwise) directly from the CMC geometry, even in this special case. Note that $\mathcal{P}_{g,k}$ does carry such a metric, and is a $K(\pi, 1)$ with $\pi = \pi_1(\mathcal{R}_{g,k})$, which when $g = 0$ is a (colored) braid group.

- Can one construct a properly (Alexandrov immersed or) embedded *minimal* surface of finite topology which is degenerate (in the sense of [22, 25]), either explicitly, or by global methods on minimal surface moduli space (for instance, might one exhibit a classifying map whose critical points must be degenerate minimal surfaces, and which is forced to have a critical point via Morse theory)? And if one works L^2 -orthogonal to the extra Jacobi fields, can one glue unduloids to the ends of the minimal surface to get a degenerate CMC surface in \mathcal{M} ? There exist minimal surfaces with all ends planar which *are* degenerate, but these are not immersed in the sense of Alexandrov; similar comments apply to the immersed constant mean curvature tori, first constructed by Wente [34], and later classified by Pinkall and Sterling [28].

- Is there a computable way to detect when an immersed surface in \mathbb{R}^3 is immersed in the sense of Alexandrov? As we have noted above, the trick of slicing with a smoothly embedded plane, and then deciding by winding number considerations that the intersection curves cannot bound an immersed domain in this plane, provides a *necessary* condition, but it would be interesting to have a general obstruction theory. The corresponding *sufficient* condition is more difficult to formulate: for example, if all such planes slice the surface in curves that bound immersed surfaces in this plane, it is tempting to think that ideas from ambient Morse theory could be used to construct the immersed 3-manifold by carefully stacking such slices. Of course, one problem is the potential non-uniqueness of the immersed surface which extends the immersed curve in each slice.

References

- [1] W. Abikoff, The real analytic theory of Teichmüller space, Springer LNM 820 (1980).
- [2] A. D. Alexandrov, Uniqueness theorems for surfaces in the large V, Vestnik Leningrad Univ. 19:13 (1958), 5–8; Amer. Math. Soc. Transl. (Ser. 2) 21 (1962), 412–416.
- [3] A. D. Alexandrov, A characteristic property of spheres, Ann. Mat. Pura Appl. 58 (1962), 303–315.
- [4] L. Bers, A remark on Mumford’s compactness theorem, Israel J. Math. 12 (1972), 400–407.
- [5] I. Biswas, On the existence of unitary flat connections over the punctured sphere with given local monodromy around the punctures, Asian J. Math. 3 (1999), 333–344.
- [6] A. Borel and A. Haefliger, La classe d’homologie fondamentale d’un espace analytique, Bull. Soc. Math. France 89 (1961), 461–513.

- [7] C. Delaunay, Sur la surface de révolution, dont la courbure moyenne est constante, *Journal de mathématiques* 6 (1841), 309–320.
- [8] J. Dorfmeister, F. Pedit, and H. Wu, Weierstrass type representation of harmonic maps into symmetric spaces, *Comm. Anal. Geom.* 6 (1998), 633–667.
- [9] T. Frankel, On the fundamental group of a compact minimal submanifold, *Annals of Math.* 83 (1966), 68–73.
- [10] K. Große-Brauckmann, New surfaces of constant mean curvature, *Math. Zeit.* 214 (1993), 527–565.
- [11] K. Große-Brauckmann, R. Kusner, and J. M. Sullivan, Constant mean curvature surfaces with three ends, *Proc. Natl. Acad. Sci. USA* 97 (2000), 14067–14068.
- [12] K. Große-Brauckmann, R. Kusner, and J. M. Sullivan, Triunduloids: Embedded constant mean curvature surfaces with three ends and genus zero, *J. Reine Angew. Math.* 564 (2003), 35–61.
- [13] K. Große-Brauckmann, R. Kusner, and J. M. Sullivan, Coplanar constant mean curvature surfaces (preprint, 2003).
- [14] D. Hoffman and W. H. Meeks, The strong halfspace theorem for minimal surfaces, *Invent. Math.* 101 (1990), 373–377.
- [15] Y. Imayoshi and M. Taniguchi, *An introduction to Teichmüller spaces*, Springer-Verlag, Tokyo, 1992.
- [16] N. Kapouleas, Complete constant mean curvature surfaces in Euclidean three-space, *Annals of Math.* 131 (1990), 239–330.
- [17] M. Kilian, I. McIntosh, and N. Schmitt, New constant mean curvature surfaces, *Experiment. Math.* 9 (2000), 595–611.
- [18] N. Korevaar and R. Kusner, The global structure of constant mean curvature surfaces, *Invent. Math.* 114 (1993), 311–332.
- [19] N. Korevaar, R. Kusner, and B. Solomon, The structure of complete embedded surfaces with constant mean curvature, *J. Diff. Geom.* 30 (1989), 465–503.
- [20] R. Kusner, R. Mazzeo, and D. Pollack, The moduli space of complete embedded constant mean curvature surfaces, *Geom. Funct. Anal.* 6 (1996), 120–137.
- [21] H. B. Lawson, The unknottedness of minimal embeddings, *Invent. Math.* 11 (1970), 183–187.
- [22] R. Mazzeo and F. Pacard, Constant mean curvature surfaces with Delaunay ends, *Comm. Anal. Geom.* 9 (2001), 169–237.
- [23] R. Mazzeo, F. Pacard, and D. Pollack, The conformal theory of Alexandrov embedded constant mean curvature surfaces, pp. 525–559 in this volume.
- [24] W. H. Meeks, The topology and geometry of embedded surfaces of constant mean curvature, *J. Diff. Geom.* 27 (1988), 539–552.
- [25] S. Montiel and A. Ros, Schrödinger operators associated to a holomorphic map, *Global differential geometry and global analysis*, Springer LNM 1481 (1991), 148–174.
- [26] D. Mumford, A remark on Mahler’s compactness theorem, *Proc. Amer. Math. Soc.* 28 (1971), 289–294.
- [27] R. Osserman, *A survey of minimal surfaces*, Van Nostrand Reinhold, New York, 1969.
- [28] U. Pinkall and I. Sterling, On the classification of constant mean curvature tori, *Annals of Math.* 130 (1989), 407–451.
- [29] J. Ratzkin, An end-to-end gluing construction for surfaces of constant mean curvature, PhD dissertation, U. Washington, 2001.
- [30] N. Schmitt, Constant mean curvature trinoids (in preparation).
- [31] M. Seppälä and T. Sorvali, *Geometry of Riemann surfaces and Teichmüller spaces*, North-Holland Mathematics Studies 169, North-Holland Publishing, Amsterdam, 1992.
- [32] K. Strebel, *Quadratic Differentials*, *Ergibnisse der Mathematik* 3.5, Springer-Verlag, Berlin, 1984.
- [33] D. Sullivan, Combinatorial invariants of analytic spaces, *Proc. of Liverpool Singularities Symp.*, Springer LNM 192 (1971), 165–169.
- [34] H. Wente, Counterexample to a conjecture of H. Hopf, *J. M. Pacificath.* 121 (1986), 193–243.

CENTER FOR GEOMETRY, ANALYSIS, NUMERICS & GRAPHICS (GANG), DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MASSACHUSETTS, AMHERST, MA 01003

E-mail address: kusner@math.umass.edu